Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.



1974 Report of

RANDOM SAMPLE EGG PRODUCTION TESTS

United States and Canada

Two-Year Combined Summary, 1972-73 and 1973-74 Range Group Rankings, 1973-74

ARS-NE-21-2

January 1975





AGRICULTURAL RESEARCH SERVICE . U.S. DEPARTMENT OF AGRICULTURE



PREFACE

Egg production tests are designed to provide poultrymen, hatcherymen, and breeders with a reliable guide to the performance of poultry stocks offered for sale. This publication contains information on many egg production traits that are of economic importance to the trade. The data were compiled from the records of official Random Sample Egg Production Tests conducted in the United States and Canada. The data resulting from these tests have been analyzed statistically by the Animal Improvement Programs Laboratory, Animal Physiology and Genetics Institute, Agricultural Research Service, USDA, Beltsville, Md.

The publication of this report is based on recommendations of the National Committee on Random Sample Poultry Testing and the Council of American Official Poultry Tests. The information was compiled by the Poultry Improvement Staff, Animal Improvement Programs Laboratory, Agricultural Research Service, from data furnished by Test supervisors.

The publication of this report does not imply approval or endorsement by the U.S. Department of Agriculture of any of the stocks mentioned.

CONTENTS

		Page
H E S D	D-year combined summary for test years 1972-73 and 1973-74	1 2 3 3 3 4 5
8	ocedures used for computing combined summary valuesStatistical methods Definition of statistical terms	12 12 13
H	nge group ranking based on 1973-74 tests	20 20 20 20
Ran	ndom Sample Egg Production Test entries and conditions, 1973-74	30
	TABLES	
1.	Two-year combined summary: Regressed means and 80% confidence limits for traits by stocks entered	6
2.	Analytical data for the traits measured, 1972-73 and 1973-74	14
3.	Factors used to adjust for test differences	15
4.	Upper and lower limits for each range group by traits and tests, 1973-74	21
5.	Range group ranking for stock entered in 1973-74 Random Sample Egg Production Tests	24
6.	Stock entered in 1973-74 tests	30
7.	Management, rations, laying house environment, and vaccination provided in tests, 1973-74	32

This report is divided into four sections:

- 1. A 2-year combined summary of the data obtained in the 1972-73 and 1973-74 Random Sample Egg Production Tests. These data were treated by acceptable statistical procedures that allow the reader to compare directly the stock entered in the various egg production tests in the United States and Canada.
- 2. An explanation of statistical procedures that were used in computing the regressed means and confidence limits of egg production traits evaluated in the 2-year combined summary.
- 3. A range group ranking for stock that was entered in 1973-74 Random Sample Egg Production Tests. The ranking shows the performance of each stock by traits compared with that of other stock in the same test.
- 4. List of stocks entered in 1973-74 tests and some of the management conditions at the test during the 1973-74 test year.

TWO-YEAR COMBINED SUMMARY FOR TEST YEARS 1972-73 and 1973-74

Entries in the various tests start with a random sample of hatching eggs or chicks of the stock to be tested. Samples are drawn according to prescribed methods to ensure that each entry is typical of the stock it represents. All entries within a test are treated alike with respect to housing, feeding, management, and disease control in order to avoid differences in performance that would be due to environment.

All tests are conducted according to these basic principles. However, even the most carefully designed and conducted tests are influenced by errors of two kinds. The first kind of error is the chance deviation or unavoidable "sampling error" made when a small sample of eggs or chicks represents an entry. The other kind of error is due to uncontrolled or unknown environmental differences between entries that occur in spite of all efforts to treat all entries within a given test as nearly alike as possible. The differences between the results for two entries in a single test for a single year may be due to these chance variations rather than to a real difference in the performance capabilities of the two stocks. The effect of such errors in comparing stocks can be materially reduced by basing comparisons on the combined results of several tests over two or more years.

If all entries compared were entered in the same tests in both years, the simple averages could be compared directly without adjustment. However, differences among tests and between years and those caused by climatic conditions and other environmental factors affect the results. As a consequence, a direct comparison of the test results of two stocks in different tests or in different years may be misleading. Therefore, to present test results in a manner that will allow sound evaluation of all stocks tested, the results were combined, by stocks and by years, and were adjusted by accepted statistical procedures for test and year differences and for variation in amount of information per stock. The results of these computations are published as the "regressed mean" for each trait for each stock that was tested (table 1).

The performance data (regressed means) reported in this summary are derived from the results reported by the individual tests for each of the past two years. It is unlikely, however, that the means for any stock, even though entered in only one test each year, will coincide precisely with the two-year average performance data as published by the test. The variations are due to adjustments for test differences, year difference, the number of tests and of years entered, and the number of replicates per test. These statistical adjustments allow predictions of what the average performance would have been for each stock had all stocks been entered in all tests each year.

The statistical treatment applied to the test data is designed to reduce the influence of nongenetic variations. This cannot be accomplished perfectly, and consequently, estimates or predictions of performance cannot be made with absolute precision. However, reliable predictions, within prescribed limitations, can be made as to whether a difference in the reported performance of stocks represents a real difference in their performance. These predictions involve the use of the confidence limit values that have been computed for each trait or performance factor reported.

A brief explanation of the statistical procedures used in computing the regressed means and confidence limits is provided in the section entitled "Procedures Used for Computing Combined Summary Values."

The following example illustrates the compilation of the two-year combined summary. This and the related explanation will help the reader to use and interpret the data in table 1.

(Illustration of regressed means and 80-percent confidence limits as they might appear for a few traits)

			FEED		EC	G G	LARG	EAND	ALBU	MEN		BLOOD	SPOTS	
STOCK CODE	BO WEI (pou	GHT	PROD PROD	GS UCED		GHT 'doz.)	EXTRA EG (per		QUAL (Haugh	.ITY	ORA	NCH MORE cent)	LESS 1/8 I (perc	NCH
CODE	RE- GRESSED MEAN	80%* CDNF. LIMITS	RE- GRESSED MEAN	80%° CDNF. LIMITS	RE- GRESSED MEAN	80%° CDNF. LIMITS	RE- GRESSED MEAN	80%* CDNF. LIMITS	RE- GRESSED MEAN	80%° CDNF. LIMITS	RE- GRESSED MEAN	80%° CDNF. LIMITS	RE- GRESSED MEAN	80%° CONF. LIMITS
-		5. 4		2.95		25.7		75.2		77.1		0.9		2.2
995	5.6	5.8	3.02	3.09	26.0	26.3	77.5	79.8	77.9	78.7	1.1	1.4	2.7	3. 2
		4.0	ĺ	2.77		25.0		69.0		80.1		0.6		0.8
996	4.2	4.4	2.83	2.89	25.2	25.4	71.0	72.8	80.9	81.7	0.7	1.0	1.1	1.4
		4.5		2.86		24.6		65.5		73.3		1.0		1.5
997	4.7	4.9	2.94	3.02	24.9	25.2	68.0	70.3	74. l	74.9	1.2	1.4	1.9	2.4
		3.7		2.73		24.9		69.2		75.5		0.9		1.2
998	4.0	4.3	2.84	2.95	25.3	25.7	72.4	75.6	76.6	77.7	1.0	1.2	1.5	1.9
		3.9		2.47		25.0		67.6		82.3		0.6		0.7
999	4.2	4.5	2.56	2.65	25.4	25.8	70.3	73.0	83.0	83.7	0.8	1.0	1.1	1.4

*If the confidence limits for two regressed means overlap, the two means are not significantly different at the 5% level.

The range of the confidence limits represents the amount of difference in the performance of two stocks that may be due to chance. If the confidence limits for two regressed means overlap, the two means are not significantly different at the 5-percent level of probability. If the confidence limits for two regressed means do not overlap, the odds are at least 19 in 20 that a real difference exists in the performance of the two stocks.

The use of the above data as a means of evaluating different stocks and traits can be illustrated as follows:

For the trait "Body Weight," the confidence limits of Stock 995 (5.4 to 5.8 lbs.) do not overlap the confidence limits of any of the other stocks. Therefore, Stock 995 has a significantly higher body weight than the others. However, the confidence limits of Stock 996 (4.0 to 4.4 lbs.) overlap the confidence limits of Stock 998 (3.7 to 4.3 lbs.) and Stock 999 (3.9 to 4.5 lbs.). The body weights of these three stocks are, therefore, not significantly different.

Using the trait "Feed per Pound of Eggs Produced" as another example, the confidence limits of Stock 995 (2.95 to 3.09 lbs.), Stock 997 (2.86 to 3.02 lbs.), and Stock 998 (2.73 to 2.95 lbs.) all overlap each other. Thus there is no significant difference in the feed conversion of these three stocks. When comparing the feed conversion of Stock 999 (2.56 lbs.) with that of the other stocks, we see that the range of its confidence limits is from 2.47 to 2.65 lbs. Since this range does not overlap the confidence limits of the other four stocks, Stock 999 has a significantly lower feed conversion than the other stocks listed.

Another example can be shown by using the trait "Albumen Quality." The confidence limits of Stock 995 (77.1 to 78.7) overlap the confidence limits of Stock 998 (75.5 to 77.7). Therefore, there is no significant difference in the albumen quality of these two stocks, even though the regressed mean of Stock 995 is 77.9 Haugh Units and Stock 998 is 76.6 Haugh Units. When Stock 995 is compared with Stocks 996 and 999, we see that the confidence limits of these two stocks do not overlap those of Stock 995. Thus, these two stocks have a significantly higher albumen quality (80.9 and 83.0 Haugh Units, respectively) than the 77.9 Haugh Units of Stock 995. In comparing Stock 995 with Stock 997, the confidence limits do not overlap. In this case, the albumen quality of Stock 997, expressed as a regressed mean of 74.1 Haugh Units is significantly lower than the regressed mean of Stock 995.

The range of the confidence limits will not necessarily be the same for two different stocks that have the same regressed mean. The number of locations in which a stock is entered, the number of replicate pens per location, the number of years entered, and the accuracy involved in adjusting for location and year effects all have a bearing on the range of the confidence limits for each individual regressed mean.

The "Income Over Feed and Chick Cost" figures reported in table 1 represent the sales value of the eggs produced and of the hens at the end of the test minus the cost of the chicks and the feed used during the growing and laying periods. These figures may be useful in comparing the overall performance of stocks, but they should not be considered as predictions of "profit" to be obtained under commercial operations. The "income" figures should be reduced by other costs, such as labor, building and equipment depreciation, vaccination, litter, interest, taxes, and insurance, to approximate profits that might be expected under commercial conditions. Surveys conducted among commercial producers indicate that such other costs may range from \$1 to \$2 per pullet housed.

Although the average chick price is reported for each stock, this value cannot be appropriately used to convert the "Income Over Feed and Chick Cost" figure to an income over feed cost figure. The average chick price shown is a simple unadjusted average of the prices reported by the entrant for his entries in the various tests and is not directly comparable to chick cost included in "Income Over Feed and Chick Cost."

Stocks Should be Compared for all Traits

All traits should be considered when using this report to evaluate the overall performance of the various stocks. The values reported for "Income Over Feed and Chick Cost" represent a composite of several traits, combined as determined by the economic conditions of the areas in which the tests are located. The conditions under which the stock is expected to perform in commercial production may differ from those prevailing at the tests, and such differences should be taken into consideration. For example, a poultryman whose local market pays unusually high premiums for large and extra large eggs should place more emphasis on egg size in his evaluation of stock than poultrymen located in areas where such premiums are not available. The local market preference for brown or white shells should also be taken into account. Traits related to interior egg quality that affect the grade are of greatest importance in areas where prices are based on quality standards.

Each person should study his local needs and conditions and then place appropriate emphasis on the performance traits that are of greatest importance to his situation. A productive and profitable stock for one poultryman under one set of conditions may not fit the needs of another poultryman under a different set of conditions.

Definition of Terms Used and Abbreviations

Stock: A term used to identify a specific breeding combination of chickens. These breeding combinations may include pure strains, strain crosses, breed crosses, incrosses, or combinations thereof. Kinds of stock and breeding methods are---

BPR Barred Plymouth Rock BX Crossbred IN White Leghorn NH New Hampshire WL PS Pure Strain Rhode Island Red White Plymouth Rock SX Strain Cross RIR WPR

Syn. Synthetic

Tests: Canada Central (C. C.) New Hampshire Cage (N.H.-C.)
Florida (Fla.) New Hampshire Floor (N.H.-F.)

Minnesota (Minn.)

Missouri Cage (Mo.-C.)

Missouri Floor (Mo.-F.)

North Carolina (N.C.)

Pennsylvania (Pa.)

Tennessee (Tenn.)

Test Year: A period beginning during the first year stated in a double-year designation and ending approximately 500 days later. See management summary shown in table 7.

	Definition of Traits
Growing mortality	Percentage of birds that died on or before the time they were 150 days old or subsequent age at housing.
Laying mortality	Percentage of birds that died after they were 150 days old or subsequent age at housing.
Age at 50 percent production	Days of age computed from the first day of the first two consecutive days of 50 percent production for living birds in the entry at that time.
Hen-housed egg production	Number of eggs laid per pullet housed computed from time of housing to the end of the test.
Hen-day egg production (to end of test)	Percent hen-day production from the time birds reached 50 percent production to end of test.
Hen-day egg production (last 30 to 60 days)	Percent hen-day production during the last 30 to 60 days of the test. Length of time involved varies according to the record keeping system of each individual test.
Feed per pound of eggs	Pounds of feed per pound of eggs produced, computed from bulk weighing of the eggs at least one day every two weeks or two days a month at equal intervals during the laying period of the test.
Egg weight	The weight of a dozen eggs computed from bulk weighing of the eggs at least one day every two weeks or two days a month during the laying period of the test.
Large and extra large eggs	Percentage of large and extra large eggs as determined by egg-size distribution computed from all eggs laid one day each week.
Albumen quality	Haugh units, computed from egg weight and albumen height of broken-out egg measured on one day's eggs per quarter, at equal intervals. The greater the Haugh units the higher the albumen quality.
Large blood spots	Percentage of ϵ_{d} gs with one or more large blood spots (1/8 inch or more in diameter), computed from at least three days' eggs per quarter, broken-out basis.
Small blood spots	Percentage of eggs with one or more small blood spots (less than 1/8 inch in diameter), computed from at least three days' eggs per quarter, broken-out basis.
Large meat spots	Percentage of eggs with one or more colored large meat spots (1/8 inch or more in diameter), computed from at least three days' eggs per quarter, broken-out basis.
Small meat spots	Percentage of eggs with one or more colored small meat spots (less than 1/8 inch in diameter), computed from at least three days' egg per quarter, broken-out basis.
Specific gravity score	Eggs are given the specific gravity score that corresponds with the specific gravity of the solution in which they will float. Eggs that do not float in 1.100 solution are given a nine score. The specific gravity of an egg is closely correlated with shell thickness; therefore, the higher the specific gravity score, the thicker the shell. Tabulation of specific gravity solutions and the corresponding specific gravity scores follow:
	Solution Score Solution Score 1.068 0 1.088 5 1.072 1 1.092 6 1.076 2 1.096 7 1.080 3 1.100 8
Podu vojeht	Average veight of hirde alive at end of test

	1:004 4	
Body weight	Average weight of birds	alive at end of test.

Income over feed and chick cost per pullet housed, with chick cost in 1,000 lots at hatch date adjusted for mortality (accidental deaths, sexing errors, and missing Income over feed and chick cost chicks not included).

Canada Central Egg Production Test

W. K. Barr, Poultry Production Section, Canada Department of Agriculture, Ottawa, Ontario, Canada Phone 613/994-9571

Florida Poultry Evaluation Center

R. B. Christmas, Chipley, Fla. 32428 Phone 904/638-0588

Minnesota Random Sample Egg Production Test

Robert E. Moehrle, Department of Agriculture, Division of Poultry Industries, 530 State Office Building, St. Paul, Minn. 55101 Phone 612/296-2861

Missouri Random Sample Egg Production Test (Cage)

Charles W. McElyea, P. O. Box 530, Mountain Grove, Mo. 65711

Phone 612/296-2861

Missouri Random Sample Egg Production Test (Floor)

Charles W. McElyea, P. O. Box 530, Mountain Grove, Mo. 65711

Phone 417/926-4151

New Hampshire Egg Production Test (Cage)

W. C. Skoglund, Department of Poultry Science, University of New Hampshire, Durham, N. H. 03824 Phone 603/862-2130

New Hampshire Egg Production Test (Floor)

W. C. Skoglund, Department of Poultry Science, University of New Hampshire, Durham, N.H. 03824 Phone 603/862-2130

North Carolina Random Sample Egg Laying Test, Salisbury

G. A. Martin, Poultry Extension Department, North Carolina State University, Raleigh, N. C. 27607 Phone 919/755-2621

Pennsylvania Random Sample Laying Test

Edgar V. Hammers, Pennsylvania Furnace, Pa. 16865

Phone 814/692-8446

Tennessee Random Sample Laying Test

H. V. Shirley, Jr., Animal Science Department, University of Tennessee, Knoxville, Tenn. 37916 Phone 615/974-7374

Copies of the final report for any of the Random Sample Egg Production Tests listed above can be obtained by writing to the test supervisor.

Table 1,--Two-year combined summary: Regressed means and 80% confidence limits for traits by stocks entered

			STOCK				MORTALITY	LITY		AGE AT	AT 50%		EGG	EGG PRODUCTION	JCTION		
STOCK	BREEDER'S NAME AND ADDRESS		STRAIN	PENS.	AVG. CHICK	GROWING	D Z	LAYING		PRODUCTION (days)	<u> </u>	HEN HOUSE	0 (10	HEN-DAY END OF TEST)		HEN-DAY (LAST 30-60 DAYS)	AY DAYS
CODE		BREEDING	OR		PRICE	(percent)	*	(percent)	*			2		(percen		(percent)	nt)
				LOCA- TIONS	(cents)	GRESSED NEAN L	80% CDNF. G	RE- GRESSED MEAN	80% CONF. G	GRESSED CC	80% CONF. GRE	GRESSED CONF.	80% RE. CONF. GRESSED LIMITS MEAN		CONF. GR	GRESSED (80%* CDNF. LIMITS
570	Animal Research Institute Ottawa, Ontario, Canada	WL PS	Kentville, R.B.C	15	30.0	3.2	2.7	4.6	7.5	176 1	172	203 2	195 211 64.	.5 66	62.5		51.1 55.5
10	Anthony, George M. & Sons Strausstown, Pa. 19559	WL SX	Anthony	57	31.9	e 6	3.8	13,1	11.3	177 1	173 181 2	14	223 68.	69 5.	67.2	ر. و و	56.8 59.8
202	Babcock Poultry Farm, Inc. Ithaca, N.Y. 14850	WL IN	Babcock B-300	114	32.8	2.5	2.0	7.6	6.4	167 1	164	245 24	237 247 74.	.3 7	73.1	8.	63.5
£443	Babcock Poultry Farm, Inc. Ithaca, N.Y. 14850	WL IN	Babcock B-303	0 m	35.0	2.7	2.4	5.1	6.8	176 1	172	31	223	9.	66.7	65.9	60.5 65.3
445	Babcock Poultry Farm, Inc. Ithaca, N.Y. 14850	RIRXSYN BX	Babcock B-380	24	32.1	2.2	1.8	6.1	7.4	180 18	176 184 2	233 24	225 241 73	17 3.	71.9	26.0	56.9 61.1
982	Canada Dept. of Agriculture Ottawa, Ontario, Canada	WL Syn.	P. D. 58	16	35.0	2.7	3.2	8.2	6.5	170 1	167 173 2	23 24	225 241 72.	2	70.3	4.5	62.2
944	Carey Farms Marion, Ohio 43302	WL SX	Carey Nick 300	4.2	32.0	2.6	2.3	5.5	4.1 7.1	176 18	172 180 2	234 24	225 243 72.	4	70.1	5.5	63.0
437	Carey Farms Marion, Ohio 43302	WL IN	Carey Nick 310	10	32.3	2.1	1.7	0.4	2.8	18	181 187 2	229 2	221 237 73.	0	71.2	£*6	67.5
289	Colonial Poultry Farm, Inc. Pleasant Hill, Mo. 64080	WL IN	True-Line 365 B	42	25.0	2.4	2.0	6.4	3.8	172 1	169 175 2	225 23	219 68	4	67.0	1.1	59.4
392	Colonial Poultry Farm, Inc. Pleasant Hill, Mo. 64080	WL IN	True-Line 365 H	9	25.0	2.6	2.3	12.5	10.4	16 173 17	169 177 2	26	217	.2 68	63.9	5.8	53,3 58,3
431	Colonial Poultry Farm, Inc. Pleasant Hill, Mo. 64080	WL IN	True-Line 365 K	13	25.0	2.5	2.1	6.6	7.9	16	167 175 22	80	67	.2 69	5.1	.1	53.6 58.6
432	Colonial Poultry Farm, Inc. Pleasant Hill, Mo. 64080	WL IN	True-Line 365 S	37	25.0	2.6	2.1	4.8	6.9	168 17	165	224 230 236	69	.9 71.	4 4	7.95	54.3 58.6
439	Colonial Poultry Farm, Inc. Pleasant Hill, Mo. 64080	RIR PS	Colonial RIR	2 1	34.0	2.5	2.3	9*6	7.9	179 18	175	193 203 213	۷,	62.	6 6	** ** **	# # # # # #

	FEED	FEED PER	EG	EGG	LARG	LARGE AND	ALBUMEN	Z		BLOOD SPOTS	SPOTS			MEAT SPOTS	POTS		0121020	0			INCOME OVER	OVER
(EGGS PRODUCED	WEI	WEIGHT	EXTRA LA EGGS	EXTRA LARGE EGGS	QUALITY	Ϋ́	1/8/1 OR N	1/8 INCH OR MORE	LESS THAN	THAN	1/8 INCH OR MORE	ICH	LESS THAN	HAN	GRAVITY) <u>}</u>	BODY	DY SHT	FEED & C	CHICK ST
STOCK		(bounds)	(02.)	(oz./doz.)	(per	(percent)	(Haugh units)	units)	(perc	(percent)	(percent)	ent)	(percent)	:nt)	(percent)	ent)	SCORE	RE	(bounds)	nds)	(dollars)	ars)
	RE- GRESSED MEAN	80%* CONF. LIMITS	RE. DRESSED MEAN	80%* CONF.	RE. GRESSED MEAN	80%* CDNF.	RE. DRESSEO MEAN	80%* CONF.	RE. GRESSED MEAN	80%* CONF.	RE. GRESSEO MEAN	80%* CONF.	RE. DRESSED MEAN	B0%* CDNF.	RE- GRESSED MEAN	80%* CDNF.	RE. DRESSEO MEAN	80%" CONF.	RE. GRESSED MEAN	80%* CONF. LIMITS	RE. GRESSEO MEAN	80%* CONF.
570	3.04	2.94	24.0	23.6	55.3	52.3 58.3	74.9	73.8	1.6	1.2	2.3	1.8	0.4	0.1	0.8	0.4	2.99	2.84	4.28	4.11	2.50	2.18 2.82
10	2.81	2.74	25.2	24.9	75.6	73.0	79.4	78.8	1.3	1.0	1.8	1.4	0.1	0.1	0.2	0.1	3.14	3.02 3.26	4-00	3.88	3.03	2.83
307	2.56	2.50	25.5	25.2	79.3	76.9	77.8	77.3	1.0	0.8	9.0	0.7	0.1	0.1	0.3	0.2	3.99	3.88	4.13	4.01	3.65	3, 32 3, 98
443	2.61	2.52	24.6	24.2	77.3	74.3	83.5	82.3 84.7	7.0	0.5	1.3	1.0	0.1	0.1	4.0	0.1	5.36	5.20	3.57	3.40	* * *	* * *
445	2.90	2.82	26.6	26.2	84.9	81.9	79.5	78.6	9.0	0.0	2.1	1.6	3.6	2.4	13.1	11.7	3.70	3.55	4.95	4.79	3.24	2.89
982	2.65	2.56	24.2	23.8	58.2	55.2	81.7	80.6	7.0	0.5	1.5	1.1	0.3	0.1	1.6	0.9	3.44	3.29	4.13	3.96	3.71	3.70
944	2.19	2.08	25.3	24.9	80.8	77.3	83.8	82.3	1.0	0.5	1.2	0.8	0.3	0.1	0.2	0.1	3.24	3.05	4.47	4.27	3.54	3.30
437	2.81	2.72	25.6	25.2	79.5	76.6	86.5	85.3	1.1	0.8	7.0	0.4	0.1	0.1	0.4	3.2	4.90	4.75	4.61	4.46	3.50	3.27 3.73
289	2.54	2.61	25.2	24.9	80.6	78.1	80.5	79.9	0.5	0.4	6*0	0.6	0.1	0.1	0.1	0.1	4.97	4.85	3.62	3.50	3.60	3,38
392	2.75	2.64	24.8	24.3	77.9	74.4	69.3	67.9	1.6	1.3	2.9	2.3	0.2	0.1	0.8	0.3	3.39	3.21	3.62	3.41	3.10	2.75
431	2.75	2.65	24.6	24.1	76.4	73.2	7.69	68.4	1.7	1.3	3.7	3.1	0.3	0.1	1.3	0.6	3.67	3.84	3.46	3.27	3.32	2.98
432	2.56	2.48	24.7	24.3	71.4	68.6	75.9	75.1	1.1	0.8	1.6	1.2	0.2	0.1	6.0	0.6	3.27	3.14	3.66	3.52	3.47	3.14
439	3.60	3.48	25.1	24.7	77.1	73.3	* * *	* * * * * *	* * *	* * * * * *	* * *	* * * * * *	* * *	* * * * * * * *	* * *	* * * * * *	* * *	* * * * * *	4.75	4.51	1.67	1.39

*If the confidence limits for two regressed means overlap, the two means are not significantly different at the 5% level.

Table 1..-Two-year combined summary: Regressed means and 80% confidence limits for traits by stocks entered (Continued)

			STOCK				MORTALITY	LITY		AGE AT 50	- 20		EGG PRO	EGG PRODUCTION	z	
STOCK	ממקים מיים מיים מיים מיים מיים מיים מיים			PENS.	AVG. CHICK	GROWING	S N C	LAYING		PRODUCTION		HEN HOUSED	HEN-DAY (TO END OF T	HEN-DAY (TO END OF TEST)	HEN-DAY (LAST 30-60 DAYS)	HEN-DAY T 30-60 DAYS)
CODE		BREEDING	STRAIN	NO.	PRICE	(percent)	nt)	(percent)	nt)	(days)	nu)	(number)	(perc	(percent)	(per	(percent)
			TRAOENAME	LOCA- TIONS	(cents)	RE- GRESSED MEAN	80%* CONF.	RE- GRESSEO MEAN	80%* CONF.	GRESSEO CONF. MEAN LIMITS	GRESSEO S MEAN	80%* CONF. LIMITS	RE- GRESSEO MEAN	80%* CONF. LIMITS	RE- GRESSEO MEAN	80%* CONF, LIMITS
339	Davis, Joe K., Hatchery Earl, N.C. 28038	RIRXBPR	Davis Combiner	38	33.5	2.9	2.4	8 . 9	7.2	1.80 1.84 1.88	196	18n 203	63.6	61.9	53.4	51.4 55.4
447	Euribrid, B. V. Boxmeer, Holland	WL SX	Hisex White	14	35.3	2.1	1.7	8.0	6.3	175 179 183	242	233	75.2	73.2	62.0	59.6
607	Fisher Poultry Farm, Ltd. Ayton, Ontario, Canada	WL SX	Fisher 107	56	32.1	2.4	1.9	10.8	9.1 12.6	173 176 179	218	212 224	68.1	66.7	57.6	55.8 59.4
99	Garber Poultry Br. Farm Modesto, Calif. 95351	WL SX	Garber G200	41	28.4	3.1	2.6	8.1	6.6 9.7	171 174 177	2.21	215 227	68.1	66.7	61.7	60.1 63.3
225	Harco Farms South Easton, Mass. 02375	RIRxBPR	Harco Sex Link	22	32.0	2.5	2.1	5.4	3.9	174 178 182	225	217	68.6	70.7	57.7	55.2 60.2
86	Hardy, C. Nelson & Son Essex, Mass. 01929	RIRXBPR BX	Deluxe Sex Link	1 6	30.0	2.0	1.7	5.4	4.0	173 177 181	207	197 217	62.2	60.09	53.7	51.2 56.2
378	Hubbard Farms, Inc. Walpole, N.H. 03608	Syn. xN. H.	Golden Comet	32	30.0	2.1	1.7	5.0	3.8	168 171 174	223	216	69.2	67.6	53.8	51.9 55.7
356	Ideal Poultry Br. Farms Cameron, Texas 76520	Syn.xWL BX	Ideal 236	71	27.0	3.2	2.6	7.9	6.6	168 171 174	230	224	70.6	69.4	63.3	61.9 64.7
234	Indiana Farm Bureau Goop. Indianapolis, Ind. 46204	WL SX	Duchess 60	37 11	32.0	3.5	2.9	12.3	10.4	169 172 175	224	218	70.7	69.3	0.09	58.3
589	Kath Line of Canada Aldergrove, B1. Col., Can.	WL SX	Kath Line H 63	16	39.0	2 + 5	2.0	11.9	9.8	169 172 175	212	204	68.3	66.4	9.69	57.3 61.º
117	Lawton Farms Foxboro, Mass. 02035	RIR×WPR BX	Buff Sex Link	1 6	32.0	2.7	2.4	3.7	2.5	173 177 181	213	204	64.0	61.7	52.6	50.1 55.1
450	Missouri Valley Hatchery Marshall, Mo. 65340	WL SX	Valley Queen	2	30.0	2.4	2.2	7.9	4.9	177 181 185	213	203 223	69.5	67.0	63.1	60.8 65.4
409	N. Cen. Reg. Plty. Br. Lab. Lafayette, Ind. 47907	WL SX	Kentville-Cornell	24	30.0	2.3	1.9	8 • 6	8.0	171 175 179	203	197	62.9	64.3	54.6	52.8 56.4

VER	X E E		80%* CONF.	1.99	3.12 3.80	2.72 3.42	2.96 3.66	3.13	2.18 2.82	2.86 3.40	3.42	3.25 3.65	2.59	2,35	2.74	2.33
INCOME OVER	FEED & CHICK COST	(dollars)	GRESSEO C	2.23 2	3.46 3	3.07 3	3.31 3	3.31	2.50	3.13	3,68	3.45	2.93	2.69	3.06	2.63
		ds)	80%* CONF. G	5.59	4.12	4.14	3.90	5.96	5.50	4.74	4.29	4.07	4.24	5.34	3.87	4.45
	WEIGHT	(spunod)	RE. GRESSEO MEAN	5.73	4.30	4.26	4.03	5.78	5.72	4.88	4.41	4.20	4.41	5.55	4.11	4.59
0	71.7 71.7		80% CONF.	2.89	4.85 5.21	3.25	4.78	3.04	3.96	3.93	4.03	3.94	2.81	3.64	* *	3.44
SPECIFIC	GRAVITY		RE. ORESSEO MEAN	3.02	5.03	3.37	4.89	3.21	3.75	3.80	4.14	4.05	2.96	3.84	* *	3.56
	HAN	cnt)	80%* CONF.	8.3	0.3	0.8	0.1	17.1	18.5	10.1	0.2	0.3	0.0	31.5	* * * * *	0.6
SPOTS	LESS THAN	(percent)	RE. GRESSEO MEAN	9.5	0.7	1.1	0.2	19.9	22.3	11.3	0.3	0.4	1.4	27.2	* * *	6.9
MEAT SPOTS	4CH ORE	ent)	BO%* CONF.	4.3	0.1	0.1	0.1	0.7	0.4	7.8	0.1	0.1	0.1	0.8	* * * * * *	0.1
	1/8 INCH OR MORE	(percent)	RE- GRESSEO MEAN	5.7	0.1	0.2	0.1	1.6	1.1	6.2	0.1	0.1	0.3	1.7	**	0.1
	NAH	ent)	80%* CONF. LIMITS	1.8	1.2	1.3	0.9	1.0	1.4	1.7	0.8	1.3	1.4	1.4	* * * * * *	1.3
SPOTS	LESS THAN 1/8 INCH	(percent)	RE. GRESSEO MEAN	2,3	1.6	1.7	1.2	1.4	1.7	2.2	1.0	1.7	1.0	1.7	**	1.7
BLOOD SPOTS	1/8 INCH OR MORE	(percent)	80%* CONF. LIMITS	1.1	0.8	1.0	0.4	0.6	0.0	1.0	5 ° 0	0.6	0.8	0.6	* * * * *	1.0
	1/8 I OR N	(per	RE- GRESSED MEAN	1.4	1.0	1.2	9.0	1.0	0.8	1.3	9.0	0.8	1.1	0.8	* *	1.3
ALBUMEN	ΥT	(Haugh units)	80%* CONF. LIMITS	75.6	78.3	74.9	81.2	80.4	74.9	77.3	75.5	81.0	75.4	76.3 79.7	* * *	75.0
ALBU	QUALITY	(Haugh	RE. ORESSEO MEAN	76.5	79.6	75.6	81.9	81.8	76.6	78.1	76.1	81.7	76.5	78.0	* *	75.8
LARGE AND	EXTRA LARGE EGGS	(percent)	80%* CONF. LIMITS	84.6	71.6	71.3	69.0	87.9	83.8	79.5	77.9	70.1	65.1	83.2	73.1	63.1
LARG	EXTRA	(ber	RE- GRESSEO MEAN	87.3	74.9	73.8	71.6	91.1	87.2	82.2	80.4	72.7	68.1	86.6	76.8	65.8
EGG	WEIGHT	(oz./doz.)	80%* CONF. LIMITS	26.3	24.9	24.7	23.9	26.1 26.9	25.4	26.5	25.2	24.6	24.5	25.3	24.6	23.8
Ē	WE	(.zo)	RE. GRESSEO MEAN	26.7	25.4	25.0	24.3	26.5	25.8	26.9	25.6	25.0	24.9	25.7	25.0	24.2
FEED PER	FGGS	(bounds)	80%* CONF. LIMITS	3.23	2.63	2.88	2.66	3.23	3.19	2.65	2.51	2.66	2.78	3.38	3.29	3.04
FEEL			RE- GRESSED MEAN	3.31	2.73	2.81	2.73	3,13	3.30	2.73	2.57	2.73	2.87	3.27	3.17	3.12
	STOCK	CODE		309	244	109	99	225	86	378	356	234	589	11.7	450	604

*If the confidence limits for two regressed means overlap, the two means are not significantly different at the 5% level.

Table 1, -- Two-year combined summary: Regressed means and 80% confidence limits for traits by stocks entered (Continued)

			STOCK	1			MORTALITY	ILITY		AGE A	T 50%		Ü	GG PRO	EGG PRODUCTION	z	
C				PENS.	AVG.	GROWING	ING.	LAYING	υz	PRODUCTION		HEN HOUSED	_	HEN-DAY	AY F TEST	HEN (LAST 30	HEN-DAY HEN-DAY
STOCK	BREEDER'S NAME AND ADDRESS	BREEDING	STRAIN	0	PRICE	(percent)	'nt)	(percent)	'nt)	(days)	5)	(number)		(percent)	ent)	(per	(percent)
			TRADENAME	LOCA- TIONS	(cents)	RE- GRESSED MEAN	80%* CONF.	RE- GRESSED MEAN	80%* CDNF.	RE- GRESSED MEAN	80%* CONF. G LIMITS	GRESSED C MEAN	80%* CONF. G LIMITS	RE. GRESSED MEAN	80%* CONF. LIMITS	RE- GRESSED MEAN	80%* CDNF. LIMITS
37	N. Cent. Reg. Plty. Br. Lab. Lafayette, Ind. 47907	WL PS	Reg. Cornell Contr.	13	30.0	2.9	3.5	12.4	10.2	179	175	215	207	6.49	65.9	54.7	52.3 57.1
352	Parks Poultry Farm Altoona, Pa. 16601	WL SX	Keystone B-1	63	33.8	3.1	2.6	6.3	5.1	174	171	233	227	72.0	70.7	60.2	58.6 61.8
382	Parks Poultry Farm Altoona, Pa. 16601	RIRXWPR BX	Sil-Go-Links	33	34.0	2.2	1.8	7.5	5.8	179	175	195 2	187	62.7	60.8	53.2	50.8 55.6
181	Shaver Poultry Br. Farm Galt, Ontario, Canada	WL SX	Starcross 288	105	31.2	2.5	2.0	5.6	6.8	173	170	245	239	75.2	73.9	64.8	63.4
451	Shaver Poultry Br. Farm Galt, Ontario, Canada	RIR SX	Starcross 579	30	35.0	3.1	3.6	8.1	6.3	173	169	211	203	66.5	64.5	54.0	51.6
999	St. Augustin Coop. Hatchery St. Augustin, Quebec, Can.	WL SX	Corvette A 1	16	31.0	2.7	2.3	11.3	9.3	176	172	216	208	69.3	67.4	63.7	61.4 66.0
401	Tatum Farms Dawsonville, Ga. 30534	WL SX	Tatum T-100	64	30.1	2.2	1.8	9.5	8.1	172	169	225	219	70.3	69.0	63.6	62.2
644	Tatum Farms Dawsonville, Ga. 30534	RlRxSyn. BX	Tatum T-173	25	30.0	1.0	0.7	5.6	4.2	171	168 174	226	234	0.69	67.3	54.3	52.1 56.5
407	Thornbers Plty. Br. Dept. Halifax, Yorkshire, Eng.	WL SX	Thornber 808	54	30.0	2.7	3.2	9.3	7.7	177	174	220	214	70.3	68.9	62.3	60.7
644	Welp's Poultry Breeding Fm. Bancroft, Iowa 50517	RIR SX	Welp Line 650 N	16	30.0	2.2	1.9	4.6	3.3	175	1 70	229	220	1.69	67.5	56.5	54°0
430	Welp's Poultry Breeding Fm. WL Bancroft, lowa 50517	WL IN	Welp Line 971	47	29.0	3.9	3.4	6.6	8.3	168	164	214	208 220	4.99	64.9	53.7	52.1 55.3
8 + + 8	Welp's Poultry Breeding Fm. WL Bancroft, Iowa 50517	WL IN	Welp Line 973	. 22	29.0	2.7	3.2	13.0	10.9	173	169	190	198	60.3	58.6 62.0	50.4	48.4

*If the confidence limits for two regressed means overlap, the two means are not significantly different at the 5% level.

Statistical Methods

The two-year combined summary includes performance data on 31 stocks that were entered in both the 1972-73 and 1973-74 tests and on 7 stocks that were entered only in the 1973-74 tests. Birds were tested at 19 locations in 1972-73 and at 21 locations in 1973-74. Table 3 lists the locations. Certain traits were not measured at some of the locations. These are identified with an NR (not reported) in the appropriate columns in table 3.

Replicate data were reported by 18 locations in 1972-73 and by 19 locations in 1973-74. The number of pens and the number of stocks tested at each location for the two years are given in table 3.

The percentage data for both years for the six traits--growing mortality, laying mortality, large blood spots, small blood spots, large meat spots, and small meat spots--were converted to angles with the arcsin transformation prior to analysis. However, the test-year adjustment factors shown in table 3 and the regressed means and confidence limits shown for these traits in table 1 are given in percent.

The replicate data were analyzed by least-squares procedures to obtain the test-year adjustment factors shown in table 3 and the repeatability estimates and the correlations among pens within tests shown in table 2. The test-year adjustment factors were then used to adjust the simple stock average for test and year effects. The adjusted stock averages (the least-squares stock means) were then regressed toward the overall mean ($^{\circ}_{\mu}$) to account for variations in number of tests entered, number of years entered, and number of replicated per test. The formula used to compute the regressed mean is:

$$\text{Regressed Mean = } \hat{\mu} + \frac{r_{2/C}}{1 + (k_3 - 1)x_1 + (k_1 - k_3)x_2 + (k_2 - k_3)r_1 + [(1/C) - k_1 - k_2 + k_3]r_2} (s)$$

where:

 \hat{x} = the average of the test and year adjusted stock means.

 r_1 = repeatability within year.

r, = repeatability from year-to-year.

 x_1 = the correlation among replicates within year and test.

 x_2 = the correlation among pens of the same stock from year-to-year for the same test.

 k_1 = an average of the number of pens per test (averaged over years).

k₂ = an average of the number of pens per year (averaged over tests).

k, = an average of the number of replicated per test-year subclass.

C = the diagonal inverse element for that stock. The reciprocal of C, i.e., $\frac{1}{C}$, is equal to nk_3 if the assumption is made that the adjustments for test-year effects are made without error; where n is the number of test-year subclasses in which that stock is entered.

s = the test-year adjusted stock average minus the overall mean $\hat{\mathbf{u}}$.

The correlations used in computing the regression coefficient were obtained from estimates of the variance components for stocks $(\hat{\sigma}_s^2)$, the stock-X-test interaction $(\hat{\sigma}_{st}^2)$, the stock-X-year interaction $(\hat{\sigma}_{sy}^2)$, and the random error $(\hat{\sigma}_e^2)$. The variance component estimates were obtained by equating the computed mean squares for these effects to their expectations. The mean squares for stocks was adjusted for the test-year subclass effects and the mean squares for the stock-X-test interaction and the stock-X-year interaction were adjusted by least-squares procedures for the effects of stocks and the test-year subclasses. The three-factor interaction was assumed to be non-existent. Ratios of the variance component estimates that were used to compute the correlations follow:

Correlation Among =
$$x_1$$
 =
$$\frac{\hat{\sigma}_s^2 + \hat{\sigma}_{st}^2 + \hat{\sigma}_{sy}^2}{\hat{\sigma}_s^2 + \hat{\sigma}_{st}^2 + \hat{\sigma}_{sy}^2 + \hat{\sigma}_{e}^2}$$
Correlations from Year-to-Year (same test) = x_2 =
$$\frac{\hat{\sigma}_s^2 + \hat{\sigma}_{st}^2 + \hat{\sigma}_{sy}^2 + \hat{\sigma}_{e}^2}{\hat{\sigma}_s^2 + \hat{\sigma}_{st}^2 + \hat{\sigma}_{sy}^2 + \hat{\sigma}_{e}^2}$$
Repeatability from Test-to-Test (within year) = r_1 =
$$\frac{\hat{\sigma}_s^2 + \hat{\sigma}_{st}^2 + \hat{\sigma}_{sy}^2 + \hat{\sigma}_{e}^2}{\hat{\sigma}_s^2 + \hat{\sigma}_{st}^2 + \hat{\sigma}_{sy}^2 + \hat{\sigma}_{e}^2}$$
Repeatability from Test-to-Test (between years) = r_2 =
$$\frac{\hat{\sigma}_s^2 + \hat{\sigma}_{st}^2 + \hat{\sigma}_{sy}^2 + \hat{\sigma}_{e}^2}{\hat{\sigma}_s^2 + \hat{\sigma}_{st}^2 + \hat{\sigma}_{sy}^2 + \hat{\sigma}_{e}^2}$$

An approximate standard error (SE) was computed for each regressed mean as follows:

SE = b
$$\sqrt{C(\hat{\sigma}_e^2 + k_1\hat{\sigma}_{st}^2 + k_2\hat{\sigma}_{sy}^2)}$$

where b is the regression coefficient given above in the formula for the regressed mean. Confidence limits were then computed for each regressed mean as follows:

Regressed Mean + 1.3 SE

The constant 1.3 was selected in order that the probability of the confidence limits overlapping by chance alone between any two means would be about 0.03. This makes the test of significance among regressed means almost comparable to using Duncan's range test at the 0.05 level of probability.

Definition of Statistical Terms

The following definitions will help the reader interpret the analytical procedures:

Overall mean	The average of the test-year adjusted means for all stocks. This is an estimate of w	what
	the overall average would have been had all stocks been entered in all tests in b	ooth
	years.	

Common stocks Stocks that are being tested at more than one location.

Test-year	The amount added to or subtracted from the actual performance of the stocks at a given
adjustment	location in a given year to bring them to the average of all the location-year sub-
factor.	classes that had complete data. These factors were determined on an intrastock basis
	with a least-squares analysis, and they are given in table 3.

	with a least-squares analysis, and they are given in table 3.
Repeatability	An intraclass correlation that measures the tendency for common stocks to rank the same
within year	from test-to-test within year. Theoretically, it can vary from 0.00 to 1.00.

relation	This correlation measures the repeatability among replicates of the same stock in the
mong	same test and year. The higher the correlation among replicates the less need
eplicates	there is for replication of stocks within test and year.

Correlation from	A correlation which measures the tendency for common stock to rank the same from year-
year-to-year	to-year when tested at the same location. The difference in the repeatability between
within tests	years and in the correlation from year-to-year within tests indicates the relative
	importance of the stock-by-test interaction.

Corr am

^{*}Includes 30 experimental stocks.

TABLE 2. --Analytical data for the traits measured 1972-73 and 1973-74

	i			Repea	tability	Correlatio	
Traits	0 "	D	,	Within	Year-to-	Among	Year-to-
	Traits Overall means Regressed means means Within year (r1) rtalitypercent- 3.5 1.9 3.9 0.1696 calitypercent- 15.3 2.9 13.7 .2093 productiondays- 12.4 164 191 .5631 egg production number- 226.2 190 253 .5314 production to end percent- 70.4 60.3 77.8 .5478 production last 30 percent- 61.0 50.4 70.4 .3530 and of eggs pounds- 2.66 2.08 3.31 .6052 ounces/dozen- 25.3 24.0 27.3 .7590 extra large eggs-percent- 76.5 55.3 91.1 .6913 ality Haugh units- 79.5 69.3 86.5 .6257 spotspercent8 .3 1.7 .1271 spots		year (r2)	replicates (X1)	year (^x 2)		
Growing mortalitypercent-	3.5	1.9	3.9	0.1696	0.0851	0.2276	0.1432
Laying mortalitypercent-	15.3	2.9	13.7	. 2093	.1761	.2864	. 2532
Age at 50% production days-	12.4	164	191	. 563 ļ	.5032	. 7232	.6632
Hen-housed egg production-number-	226.2	190	253	.5314	. 4855	.6447	. 5988
Hen-day egg production to end of testpercent-	70.4	60.3	77.8	. 5478	.5100	. 6650	. 6271
Hen-day egg production last 30 to 60 dayspercent-	61.0	50.4	70.4	. 3530	.3367	. 4991	.4828
Feed per pound of eggs pounds-	2.66	2.08	3,31	.6052	.5608	. 7590	. 7146
Egg weight ounces/dozen-	25.3	24.0	27.3	. 7590	.6439	. 8453	.7301
Large and extra large eggs-percent-	76.5	55.3	91.1	.6913	. 5741	.8050	.6878
Albumen quality Haugh units-	79.5	69.3	86.5	.6257	.6242	.6727	.6712
Large blood spotspercent-	. 8	. 3	1.7	.1271	.1151	.2385	. 2265
Small blood spotspercent-	1.4	. 7	4.1	.1246	.1212	. 2527	.2494
Large meat spotspercent-	. 5	. 1	5.7	.7144	.6644	.8144	. 7644
Small meat spotspercent-	1.3	.0	27.2	.8176	.8149	.8635	.8609
Specific gravity score-	4.0	1.87	5.72	. 6373	. 5908	.6783	. 6318
Body weight pounds-	4.45	3.46	5.88	.8418	.8008	.9308	.8898
Income over feed and chick costdollars-	3.28	1.67	4.96	. 4650	.4350	. 5800	. 5500

NOTE: The values for these factors are based on the 38 commercially available stocks as well as the 30 experimental stocks that were tested. The individual performance data for the experimental entries were analyzed but not published in this report.

TABLE 3. -- Factors used to adjust for test differences

	Pe	ns	Stocks	tested	Mortality (percent)									
Test	(num		(num		Growing period		g period							
	1973	1974	1973	1974	1973 1974	1973	1974							
Central Canada No. 6 - (2/cage)	48	48	12	12	-3.50 +01.83	+3.71	+2.29							
Central Canada No. 7 - (2/cage)	48	48	12	12	+3.95 + 2.96	- 3.20	- 3.01							
Florida No 1 - Floor		24		12	÷ 2.69	ter day	÷1.38							
Florida No. 2 - (2/cage)	48	48	12	12	- .94 + 1.21	74	+1.89							
Florida No. 6 - Floor		24	e =	12	÷ 3.74	~ =	+3.34							
Florida No. 5 - (2/cage)	48	48	12	12	+ .13 + .50	-1.31	01							
Minnesota No. 1 - Floor	14	10	14	10	-1.00 + 4 38	- .87	÷ .98							
Minnesota No. 4 = (3/cage)	39	33	13	11	- 3 43 + 4.05	- 3.08	÷1.74							
Missouri Cage - (8/cage)	111	28	16	14	÷1.31 + 28	+8.22	+7.70							
Missouri Floor	88	54	22	27	+1.34 + .28	+2.96	-1.75							
New Hampshire No. 7 - (3/cage)	120	135	15	17	+3.79 + 2.62	~1.06	+ .86							
New Hampshire No. 4 - Floor	24	24	8	8	+3.12 + .91	-3.00	-1.7 1							
North Carolina No. 3 - Floor	40	20	20	10	+2.68 + 1.32	+3.70	+5.34							
North Carolina No. 4 - (2/cage)		40	20	10	+3.78 + 2.08	12	+1.39							
North Carolina No. 5 - (7/cage)	40	20	20	10	+3.49 - 1.78	-1.94	+1.68							
Pennsylvania No. 1 - Floor	48	48	24	24	+ .04 + 1.78	+3.66	-1,45							
Pennsylvania No. 2 - (3/cage)	48	48	24	24	-2.07 + 1.36	-8.41	- 7.51							
Tennessee No. 5 - (2/cage)	26	28	13	14	+ .11 - 5.92	10	+ .22							
Tennessee No. 6 - (2/cage)	26	28	13	14	38 - 6.40	+4.73	+ .73							
Tennessee No. 7 - (2/cage)	26	28	13	14	+ .04 - 5.99	-3.39	97							
Tennessee No. 8 - (2/cage)	26	28	13	14	+ .04 -17.89	- 5.43	-2.78							

 ${\tt TABLE~3.\text{--}Factors~used~to~adjust~for~test~differences\text{--}Continued}$

	50 pe	e at rcent			1	-day	Hen-day				
Test		iction ys)		noused nber)	l '	l of test) cent)	(last 30-6 (perc	, ,			
	1973	1974	1973	1974	1973	1974	1973	1974			
Central Canada No. 6 - (2/cage)	+ 6.19	+ 2.51	-10.23	+ 1.21	-2.16	+ 0.91	- 0.40	- 3.04			
Central Canada No. 7 - (2/cage)	+ .62	+15.05	+ 2.44	+ 2.36	-1.07	+ 1.47	- 2.59	- 4.36			
Florida No. 1 - Floor		- 4.97		- 5.20		- 2.15	e- to	+ 4.13			
Florida No. 2 - (2/cage)	- 2.99	- 5.20	+ .61	- 6.13	-1.70	- 3.05	- 4.11	+ 1.16			
Florida No. 6 - Floor		- 3.08		- 9.52		+ .09		- 3,34			
Florida No. 5 - (2/cage)	+ 4.97	- 6.50	+ 3.24	+11.21	-2.57	+ .34	+ 3.81	+ 3.98			
Minnesota No. 1 - Floor	+ 7.20	- 2.74	- 4.70	- 6.79	-1.30	- 2.49	+ 4.16	- 2.36			
Minnesota No. 4 - (3/cage)	+ 5.95	- 7.18	+ 4.04	+ 4.84	+ .23	+ .14	+ 5.45	64			
Missouri Cage - (8/cage)	- 8.45	+ 6.43	+ 5.74	-10.99	-2.24	- 1.34	- 4.57	NR∗			
Missouri Floor	-10.75	+ 2.33	+ 6.85	-10.54	+1.43	- 1.98	76	NR*			
New Hampshire No. 7 - (3/cage)	+ 8.83	+ 7.84	- 7.58	+20.11	-1.75	+ 7.71	NR*	- 1.68			
New Hampshire No. 4 - Floor	- 3.80	+10.05	+ 3.87	- 1.06	-1.81	+ .04	NR*	+ .62			
North Carolina No. 3 - Floor	84	-18.38	-18.01	+ .05	-2.10	+ 6.22	- 2.36	- 2.71			
North Carolina No. 4 - (2/cage)	+ 3.88	+ .34	-22.52	-24.39	-6.01	- 6.56	- 5.39	- 5.67			
North Carolina No. 5 - (7/cage)	+ 9.38	+ .97	+ .82	+ 7.69	+1.02	+ 3.20	- 3.38	- 6.45			
Pennsylvania No. 1 - Floor	12	- 4.68	-15.20	- 4.70	-2.78	- 2.19	+ 1.41	- 6.99			
Pennsylvania No. 2 - (3/cage)	- 3.68	- 6.57	+17.20	+ 8.79	+1.35	69	+12.39	+10.04			
Tennessee No. 5 - (2/cage)	-14.59	-15.71	- 7.32	+ 9.60	-2.83	+ 2.71	80	- 1.91			
Tennessee No. 6 - (2/cage)	- 7.07	- 8.19	- 8.50	+ 4.90	-1.25	+ 1.86	+ 2.97	+ 1.40			
Tennessee No. 7 - (2/cage)	+ 4.93	+ 3.81	- 7.65	+11.89	-2.13	+ 4.64	- 5.09	- 4.48			
Tennessee No. 8 - (2/cage)	+ 4.93	+21.26	+ 7.16	+31.75	+1.60	+15.08	+ 7.36	+ 8.57			

^{*} Data for this trait not reported.

TABLE 3. -- Factors used to adjust for test differences -- Continued

Test	of e	er pound eggs inds)	Egg w	ozen)	large (per	e eggs	(Haugh	n quality
Central Canada No. 6 - (2/cage)	+0,32	+0.23	1973 -0.58	1974	1973 -06.27	-09.95	+00.61	<u>1974</u> -3.65
Central Canada No. 7 - (2/cage)	+ .06	+ .24	- .63	+1.05	69	+28.02	02	+ .93
Florida No. 1 - Floor		+ .22		93		- 8.92		-2.04
Florida No. 2 - (2/cage)	+ .25	+ .27	-1.37	-1.24	-15.26	-10.99	+ 7.02	+3.74
Florida No. 6 - Floor		03		04		- 2.23		-2.31
Florida No. 5 - (2/cage)	15	23	+ .11	15	+10.58	+ 4.48	÷ 1.61	~2.96
Minnesota No. 1 - Floor	10	+ .28	+1.00	82	+ 4.08	-12.09	+ 2.51	-4.17
Minnesota No. 4 - (3/cage)	21	40	43	82	+ 5.75	+ .55	48	-1.77
Missouri Cage - (8/cage)	+ .01	+ .41	67	+ .01	-10.10	- 4.40	-13.32	-8.79
Missouri Floor	06	+ .27	21	+ .31	- 4.03	- 2.58	-10.33	NR*
New Hampshire No. 7 - (3/cage)	+ .08	35	+ .17	+ .78	+14.75	+12.46	+ 3.57	+1.78
New Hampshire No. 4 - Floor	+ .04	+ .23	72	+1.03	- 2.50	+28.47	54	+ .03
North Carolina No. 3 - Floor	03	04	- .36	55	-15.38	-15.33	- 1.89	-1.76
North Carolina No. 4 - (2/cage)	+ .31	+ .29	19	54	-10.20	- 6.66	+ 5.45	+2.91
North Carolina No. 5 - (7/cage)	04	29	+ .61	+ .95	+15.48	+10.45	+ 2.92	17
Pennsylvania No. 1 - Floor	06	48	+ .68	+ .41	- 7.08	- 4.46	- 2.62	+ .12
Pennsylvania No. 2 - (3/cage)	+ .04	+ .13	97	-1.29	-14.99	-11.24	+ 4.84	+3.40
Tennessee No. 5 - (2/cage)	+ .04	+ .03	31	44	- 9.92	-11.71	- 1.42	-4.82
Tennessee No. 6 - (2/cage)	+ .22	+ .29	67	97	- 9.20	-13.30	- 2.26	-3.95
Tennessee No. 7 - (2/cage)	15	19	+ .94	+ .79	+11.15	+ 6.93	+ 3.81	+1.43
Tennessee No. 8 - (2/cage)	27	99	+1.04	+4.65	+10.11	+45.01	+ .68	+7.23

^{*} Data for this trait not reported.

TABLE 3. -- Factors used to adjust for test differences--Continued

	Blood	dspots	Bloo	d spots	Meat	spots	Meat spots				
Test	i	or more	1	n 1/8 inch	1/8 inch			n 1/8 inch			
	1973	1974	(per 1973	1974	(perc-	ent) 1974	(per	1974			
Central Canada No. 6 - (2/cage)	+0.08	-01.21	+1.39	+0.74	+1.15	+1.04		+03.16			
Central Canada No. 7 - (2/cage)	-3.35	- 1.70	-1.82	-2.21	-2.44	-2.45	- 3.51	- 3.25			
Florida No. 1 - Floor		- 1.87		+2.48		-1.14		- 1.63			
Florida No. 2 - (2/cage)	-2.81	- 2.64	-1.81	52	-1.41	-3.52	+ 1.54	14			
Florida No. 6 - Floor		46		07		+ .28		+ .25			
Florida No. 5 - (2/cage)	+ .28	- 1.80	+1.00	+ .54	+2.71	+2.34	+ 2.04	+ 1.85			
Minnesota No. 1 - Floor	+3.26	- 2.75	+ .02	+4.02	+2.15	+ .67	+ 5.05	+ 1.32			
Minnesota No. 4 - (3/cage)	03	+ .99	+ .05	+1.15	+2,52	+2.54	+ 3.07	+ 1.37			
Missouri Cage - (8/cage)	+1.45	NR*	+3.25	NR*	+1.38	ÑR≉	+ 2.99	NR*			
Missouri Floor	+1.52	NR*	+4.05	NR*	+1.50	NR*	+ 3.63	NR*			
New Hampshire No. 7 - (3/cage)	+2.39	+ 2.89	+6.93	+5.96	+1.34	-1.02	-11.03	- 7.76			
New Hampshire No. 4 - Floor	-3.84	- 2.06	-1.89	-1.35	-2.45	-1.76	- 3.60	- 2.80			
North Carolina No. 3 - Floor	-1.79	- 2.33	- 3.89	-3.30	69	26	+ .80	+ .82			
North Carolina No. 4 - (2/cage)	-2.22	- 1.39	71	+ .51	-2.00	-3.61	+ .98	+ .31			
North Carolina No. 5 - (7/cage)	+2.31	+ 1.95	+2.52	+7.36	+2.62	55	-11.62	-12.34			
Pennsylvania No. 1 - Floor	+ .37	+ .07	-5.84	-2.04	+ .28	+ .30	03	+ 1.26			
Pennsylvania No. 2 - (3/cage)	-2.67	- 1.45	61	+ .31	-1.03	-4.19	+ 1.78	+ .06			
Tennessee No. 5 - (2/cage)	-1.16	- 1.11	-1.89	07	+1.30	+ .43	+ 3.32	+ 3.06			
Tennessee No. 6 - (2/cage)	 13	+ 1.14	+1.88	13	+ .84	+1.24	+ 2.90	+ 3.61			
Tennessee No. 7 - (2/cage)	+2.93	+ 2.74	-3.49	+3.27	+ .38	+ .34	+ .71	+ 1.49			
Tennessee No. 8 - (2/cage)	+2.88	+12.16	+ .70	- 6.78	+ .63	+2.57	+ 2.90	+ 9.59			

^{*} Data for this trait not reported.

TABLE 3. -- Factors used to adjust for test differences -- Continued

Test	sc	ic gravity	(pou	weight	feed chick (doll	and cost
	1973	1974	1973	1974	1973	1974
Central Canada No. 6 - (2/cage)	-1.08	-1.85	+0.25	+0.10	+1.56	+0.80
Central Canada No. 7 - (2/cage)	+ .74	+1.02	23	+ .22	+1.57	+ .68
Florida No. 1 - Floor	- ~	07		11		NR*
Florida No. 2 - (2/cage)	+1.65	+1.14	25	08	NR*	NR*
Florida No. 6 - Floor	-	26	00 ftg	07	***	NR*
Florida No. 5 - (2/case)	64	-1.65	+ .02	16	NR*	NR*
Minnesota No. 1 - Floor	~ .91	-1.66	+ .02	+ .29	+ .35	+1.73
Minnesota No. 4 - (3/cage)	84	-1.33	+ .04	24	+ .56	+1.80
Missouri Cage - (8/cage)	86	39	42	13	22	NR*
Missouri Floor	48	NR*	34	18	-1.21	39
New Hampshire No. 7 - (3/cage)	+ .89	+ .58	21	+ .02	-2.14	-1.33
New Hampshire No. 4 - Floor	+ .85	+1.09	16	+ .28	-2.82	71
North Carolina No. 3 - Floor	+ .21	09	04	10	+ .56	+ .03
North Carolina No. 4 - (2/cage)	+1.20	+1.15	19	+ .04	+1.03	+ .44
North Carolina No. 5 - (7/cage)	+1.38	+ .81	12	08	+1.68	+ .95
Pennsylvania No. 1 - Floor	+ .23	+ .59	+ .11	+ .23	-2.08	27
Pennsylvania No. 2 - (3/cage)	+1.35	+1.11	02	+ .18	-1.93	92
Tennessee No. 5 - (2/cage)	-1.28	-1.41	03	+ .30	+ .87	90
Tennessee No. 6 - (2/cage)	-1.55	-1.29	+ .26	+ .34	+ .97	86
Tennessee No. 7 - (2/cage)	+ .28	01	14	+ .22	+ .74	82
Tennessee No. 8 - (2/cage)	+ .09	+ .49	13	+ .33	+1.02	92

^{*} Data for this trait not reported.

RANGE GROUP RANKING BASED ON 1973-74 TESTS

How Group Rankings Were Determined for Each Trait

The information in this section deals only with the test data obtained during the 1973-74 test year.

The performance of each entry in the 10 Random Sample Egg Production Tests conducted during 1973-74 is reported as the Range Group Rank of the entry for the trait measured. These rankings were determined in the following manner. For each trait the entries in each test were alined in descending order of performance from the most desirable to the least desirable. The "mean" or average performance for the trait was then determined. All entries above the mean are in range group 1 or 2, and those below the mean are in range group 3 or 4. The dividing point for the entries above or below the mean is the midpoint of the range between the mean and the top or bottom entry. An illustration follows.

Stocks entered in the Pennsylvania test had a mean, or average, of 224.40 eggs for the trait "Egg Production per Hen Housed." The highest average number of eggs laid by an entry in this test was 251.20 and the lowest average number laid by an entry was 170.70 eggs. To arrive at the dividing point between the first and second range groups, the mean (224.40) was subtracted from the highest number of eggs (251.20). The result, 26.80 eggs, was divided by two to get the midpoint of the range (13.40 eggs). This was then subtracted from the highest number of eggs (251.20 minus 13.40) to arrive at the dividing point (237.80 eggs) between the first and second range groups. To determine the dividing point between the third and fourth range groups, the same procedure was used, except that the lowest average number of eggs (170.70) was subtracted from the mean (224.40) This difference, or range, (53.70 eggs) was then divided by two, and the result (26.85 eggs) was subtracted from the mean (224.40 minus 26.85) to get the dividing point (197.55 eggs) between the third and fourth range groups. These determinations for ten traits from each test are tabulated in table 4.

The breeders of the stock tested and the Range Group Ranking, by traits, of each entry of the stock are shown in table 5. Each entry is also identified by the abbreviated name of the entrant. If the sample was drawn from a source other than the entrant's hatchery or supply flock, the abbreviated name of the source of the sample is shown in parentheses following the entrant's name.

The listing of the entries in the four range groups, with all entries of each stock in one table, allows the reader to quickly evaluate a stock based on this method of analysis. It should be kept in mind, however, that this method provides just four broad classifications. One-tenth of an egg or one-tenth of a percent difference in mortality could move an entry up or down one Range Group Rank, depending on its place in the range grouping.

Tabular Listing of Stock Entered in Tests

The listing of all stock entered in the 1973-74 Random Sample Egg Production Tests is given in table 6. This listing will permit the reader to see at a glance the abbreviated name of the breeder of the stock, the strain or trade name of the stock, and the total number of entries of each stock which were tested during 1973-74. The tests in which each stock was entered are also given.

Management and Environmental Conditions at Tests

Some of the more important management and environmental conditions found in the individual tests during the 1973-74 testing year are found in table 7. Other conditions at the various testing stations were undoubtedly different. However, the important consideration is that all entries at a given location were treated as nearly alike as possible.

			Tests	
Traits measured	Central			Missouri
	Canada	Florida	Minnesota	Cage
Income over feed and chick cost;	2 420		1.731	
Averagedol./hen housed-	2.430 3.270 - 2.850		2.370 - 2.050	
Range group 1	2.849 - 2.430	Not Reported	2.049 - 1.731	Not Reported
Range group 2	2.429 - 1.725	Not reported	1.730 - 1.390	not iteported
Range group 4	1. 724 - 1. 020		1.389 - 1.050	
Egg production;	1.724 - 1.020		1. 307 - 1.030	
Average number/hen housed-	223.17	228.18	239.02	222,96
Range group 1		248.60 - 238.39	264.20 - 251.61	247.00 - 234.98
Range group 2		238.38 - 228.18	251.60 - 239.02	234.97 - 222.96
Range group 3		228.17 - 206.34	239.01 - 229.01	222.95 - 208.08
Range group 4		206.33 - 184.50	229.00 - 219.00	208.07 - 193.20
Age at 50 percent production;				
Averagedays-	159.6	177.3	168.2	191.6
Range group 1	156.0 - 157.8	172.0 - 174.7	162.0 - 165.1	183.0 - 187.3
Range group 2	157.9 - 159.6	174.8 - 177.3	165.2 - 168.2	187.4 - 191.6
Range group 3	159.7 - 161.8	177.4 - 179.7	168.3 - 175.1	191.7 - 196.8
Range group 4	161.9 - 164.0	179.8 - 182.0	175.2 - 182.0	196.9 - 202.0
Growing mortality;				
Averagepercent-	3,22	3.06	2.29	2.12
Range group 1	1.40 - 2.31	1.70 - 2.38	1.00 - 1.65	0 1.06
Range group 2	2.32 - 3.22	2.39 - 3.06	1.66 - 2.29	1.07 - 2.12
Range group 3	3.23 - 5.11	3.07 - 4.73	2.30 - 3.15	2.13 - 3.31
Range group 4	5.12 - 7.00	4.74 - 6.40	3.16 - 4.00	3.32 - 4.50
Laying mortality;	11 71	7 22	4 02	r 71
Averagepercent-	11.71	7.33	4.92	5.71
Range group 1	6.10 - 8.90	3.30 - 5.32	1.40 - 3.16	1.30 - 3.51
Range group 2	8.91 - 11.71	5.33 - 7.33	3.17 - 4.92	3.52 - 5.71
Range group 4	11.72 - 15.40 15.41 - 19.10	7.34 - 11.47 11.48 - 15.60	4.93 - 8.51 8.52 - 12.10	5.72 - 11.01 11.02 - 16.30
Egg weight;	13.41 - 17.10	11.40 - 15.00	0.52 - 12.10	11.02 - 10.30
Averageounces/dozen-	24.18	25.66	25.32	25.64
Range group 1	26.20 - 25.19	26.40 - 26.03	26.30 - 25.81	27.30 - 26.47
Range group 2	25.18 - 24.18	26.02 - 25.66	25.80 - 25.32	26.46 - 25.64
Range group 3	24.17 - 23.69	25.65 - 25.13	25.31 - 24.76	25.63 - 24.97
Range group 4	23.68 - 23.20	25.12 - 24.60	24.75 - 24.20	24.96 - 24.30
Large and extra large eggs;				
Averagepercent-	47.27	86.15	81.65	89.95
Range group I	72.10 - 59.68	91.30 - 88.73	86.30 - 83.98	97.40 - 93.68
Range group 2	59.67 - 47.27	88.72 - 86.15	83.97 - 81.65	93.67 - 89.95
Range group 3	47.26 - 40.13	86.14 - 82.28	81.64 - 76.63	89.94 - 82.38
Range group 4	40.12 - 33.00	82.27 - 78.40	76.62 - 71.60	82.37 - 74.80
Feed per pound of eggs;	2 402	2 402	2 2/2	2 722
Averagepounds-	2.483	2.483	2.262	2.722
Range group 1	2.320 - 2.402	2.330 - 2.406	2.080 - 2.171	2.500 - 2.611
Range group 2	2.403 - 2.483	2.407 - 2.483	2.172 - 2.262	2.612 - 2.722
Range group 3	2.484 - 2.677 2.678 - 2.870	2.484 - 2.596	2.263 - 2.341	2.723 - 2.981
	2.010 - 2.010	2.597 - 2.710	2.342 - 2.420	2.982 - 3.240
Albumen quality; AverageHaugh units-	77.75	82.76	87.54	
Range group 1	81.80 - 79.78	84.70 - 83.73	90.50 - 89.02	
Range group 2	79.77 - 77.75	83.72 - 82.76	89.01 - 87.54	Not Reported
Range group 3	77.74 - 76.23	82.75 - 81.43	87.53 - 86.17	2.07 Ltoported
Range group 4	76.22 - 74.70	81.42 - 80.10	86.16 - 84.80	
Blood spots, all sizes;				
Averagepercent-	4.31	2.88	1.69	
Range group 1	1.60 - 2.95	1.30 - 2.09	0 - 0.85	
Range group 2	2.96 - 4.31	2.10 - 2.88	.86 - 1.69	Not Reported
Range group 3	4.32 - 8.20	2.89 - 4.24	1.70 - 3.25	•
Range group 4	8.21 - 12.10	4.25 - 5.60	3.26 - 4.80	

TABLE 4.--Upper and lower limits for each range group by traits and tests, 1973-74--(Continued)

		Tests	
Traits measured	Missouri	New Hampshire	New Hampshire
	Floor	Cage	Floor
Income over feed and chick cost;			
Averagedol./hen housed-	3.501	4.713	3.596
Range group 1	5.580 - 4.541	5.730 - 5.221	5.390 - 4.493
Range group 2	4.540 - 3.501	5.220 - 4.713	4.492 - 3.596
Range group 3	3.500 - 2.406	4.712 - 4.071	3.595 - 2.838
Range group 4	2.405 - 1.310	4.070 - 3.430	2.837 - 2.080
Egg production;			
Averagenumber/hen housed-	224.37	222.48	199.33
Range group 1	270.10 - 247.24	246.80 - 234.64	236.10 - 217.71
Range group 2	247.23 - 224.37	234.63 - 222.48	217.70 - 199.33
Range group 3	224.36 - 206.69	222.47 - 209.54	199.32 - 186.11
Range group 4	206.68 - 189.00	209.53 - 196.60	186.10 - 172.90
Age at 50 percent production;			
Averagedays-	178.7	173.9	171.0
Range group 1	169.0 - 173.9	157.0 - 165.4	152.0 - 161.5
Range group 2	174.0 - 178.7	165.5 - 173.9	161.6 - 171.0
Range group 3	178.8 - 183.4	174.0 - 185.4	171.1 - 175.5
Range group 4	183.5 - 188.0	185.5 - 197.0	175.6 - 180.0
Growing mortality;			
Averagepercent-	1.66	3.52	1.75
Range group 1	.30 - 0.98	0 - 1.76	0 - 0.88
Range group 2	.99 - 1.66	1.77 - 3.52	.89 - 1.75
Range group 3	1.67 - 2.68	3.53 - 8.91	1.76 - 3.88
Range group 4	2.69 - 3.70	8.92 - 14.30	3.89 - 6.00
Laying mortality;			
Averagepercent-	11.35	6.12	5.68
Range group 1	2.50 - 6.92	1.60 - 3.86	2.20 - 3.94
Range group 2	6.93 - 11.35	3.87 - 6.12	3.95 - 5.68
Range group 3	11.36 - 17.57	6.13 - 9.31	5.69 - 10.04
Range group 4	17.58 - 23.80	9.32 - 12.50	10.05 - 14.40
Egg weight;			
Averageounces/dozen-	25.04	24.93	25.68
Range group 1	27.40 - 26.22	26,50 - 25,71	26.50 - 26.09
Range group 2	26.21 - 25.04	25.70 - 24.93	26.08 - 25.68
Range group 3	25.03 - 24.17	24.92 - 24.46	25.67 - 25.29
Range group 4	24.16 - 23.30	24.45 - 24.00	25.28 - 24.90
Large and extra large eggs;			
Averagepercent-	82.07	69.61	75.61
Range group 1	94.40 - 88.24	87.70 - 78.66	83.40 - 79.51
Range group 2	88.23 - 82.07	78.65 - 69.61	79.50 - 75.61
Range group 3	82.06 - 72.99	69.60 - 62.21	'75.60 - 69.16
Range group 4	72. 98 - 63. 90	62.20 - 54.80	69. 15 - 62. 70
Feed per pound of eggs;			
Averagepounds-	3.249	3.028	3.335
Range group 1	2.810 - 3.029	2.660 - 2.843	2.680 - 3.008
Range group 2	3.030 - 3.249	2.844 - 3.028	3.009 - 3.335
Range group 3	3.250 - 3.659	3.029 - 3.149	3.336 - 3.598
Range group 4	3.660 - 4.070	3.150 - 3.270	3.599 - 3.860
Albumen quality;			
Average Haugh units-		79.74	78.29
Range group 1		84.60 - 82.17	82.70 - 80.49
Range group 2	Not Reported	82.16 - 79.74	80.48 - 78.29
Range group 3		79.73 - 78.07	78.28 - 76.64
Range group 4		78.06 - 76.40	76.63 - 75.00
Blood spots, all sizes;			
Averagepercent-		0.99	2.36
Range group 1		0 - 0.49	0 - 1.18
Range group 2	Not Reported	.5099	1.19 - 2.36
Range group 3		1.00 - 2.59	2.37 - 5.38
Range group 4		2.60 - 4.20	5.39 - 8.40

TABLE 4. -- Upper and lower limits for each range group by traits and tests, 1973-74-- (Continued)

		Tests	
Traits measured	North		
	Carolina	Pennsylvania	Tennessee
Income over feed and chick cost;	0.510		
Average dol./hen housed-	2.710	3, 908	4.069
Range group 1	3.520 - 3.115	5,550 - 4,729	4.790 - 4.430
Range group 2	3.114 - 2.710	4.728 - 3.908	4.429 - 4.069
Range group 3	2.709 - 2.205	3.907 - 2.774	4.068 - 3.420
Range group 4	2.204 - 1.700	2.773 - 1.640	3.419 - 2.770
Egg production; Average number/hen housed-	229.18	224,40	208.77
Range group 1	258.50 - 243.84	251.20 - 237.80	228.00 - 218.39
Range group 2	243.83 - 229.18	237.79 - 224.40	218.38 - 208.77
Range group 3	229.17 - 214.99	224. 39 - 197. 55	208.76 - 188.99
Range group 4	214.98 - 200.80	197.54 - 170.70	188.98 - 169.20
Age at 50 percent production;			
Averagedays-	177.1	181.0	168.2
Range group 1	166.0 - 171.6	173.0 - 177.0	165.0 - 166.6
Range group 2	171.7 - 177.1	177.1 - 181.0	166.7 - 168.2
Range group 3	177.2 - 187.1	181.1 - 187.5	168.3 - 172.6
Range group 4	187.2 - 197.0	187.6 - 194.0	172.7 - 177.0
Growing mortality;			
Average percent-	2.28	0.93	7. 75
Range group 1	1.10 - 1.69	0 - 0.47	0 - 3.88
Range group 2	1.70 - 2.28	.4893	3.89 - 7.75
Range group 3	2.29 - 3.39	. 94 - 1.67	7.76 - 11.38
Range group 4	3.40 - 4.50	1.68 - 2.40	11.39 - 15.00
Laying mortality;	10 55	4 57	11 20
Average percent-	10.55 5.20 - 7.88	6.57 1.50 - 4.03	11.30 5.00 - 8.15
Range group 1Range group 2	7.89 - 10.55	4.04 - 6.57	8. 16 - 11. 30
Range group 3	10.56 - 14.68	6.58 - 10.68	11.31 - 14.00
Range group 4	14.69 - 18.80	10.69 - 14.80	14.01 - 16.70
Egg weight;	11,07 - 10,00	10,07 - 11,00	11,01 - 10,10
Averageounces/dozen-	26.49	26.18	24.51
Range group 1	27.80 - 27.15	28.80 - 27.49	26.00 - 25.25
Range group 2	27.14 - 26.49	27.48 - 26.18	25.24 - 24.51
Range group 3	26.48 - 25.50	26.17 - 25.24	24.50 - 23.75
Range group 4	25.49 - 24.50	25.23 - 24.30	23.74 - 23.00
Large and extra large eggs;			
Average percent-	88.27	77.12	69.71
Range group 1	95.80 - 92.04	90.00 - 83.56	82.10 - 75.91
Range group 2	92.03 - 88.27	83. 55 - 77. 12	75.90 - 69.71
Range group 3	88.26 - 83.19	77.11 - 65.81	69.70 - 60.16
Range group 4Feed per pound of eggs;	83.18 - 78.10	65, 80 - 54, 50	60.15 - 50.60
Averagepounds-	2.573	3 015	2 014
Range group 1	2.320 - 2.447	3.015 2.560 - 2.788	2.914
Range group 2	2.448 - 2.573	2.789 - 3.015	2.730 - 2.822 2.823 - 2.914
Range group 3	2.574 - 2.832	3.016 - 3.223	2.915 - 3.127
Range group 4	2.833 - 3.090	3.224 - 3.430	3. 128 - 3. 340
Albumen quality;	2,000 - 0,070	J. 221 - J. TJU	J. 120 - J. J. U
Average Haugh units-	76.13	82.43	78.58
Range group l	79.10 - 77.62	86.40 - 84.41	85.00 - 81.79
Range group 2	77.61 - 76.13	84.40 - 82.43	81.78 - 78.58
Range group 3	76.12 - 74.72	82.42 - 80.01	78.57 - 76.59
Range group 4	74.71 - 73.30	80.00 - 77.60	76.58 - 74.60
Blood spots, all sizes;			
Average percent-	3. 36	3.19	4.64
Range group 1	1.50 - 2.43	1.40 - 2.29	1.30 - 2.97
Range group 2	2.44 - 3.36	2.30 - 3.19	2.98 - 4.64
Range group 3	3.37 - 4.33	3.20 - 4.89	4.65 - 9.82
Range group 4	4.34 - 5.30	4.90 - 6.60	9.83 - 15.00

TABLE 5. --Range group ranking for stock entered in 1973-74 random sample egg production tests

STO98	8	1	•	2 3		3		٠,		3	4	3	2	П		1		ı	_	_	3		7		-
MENMENA #,	4	1	1	2 %		2	4.	4		4	4	3	3	4		1		1	2	1	2		_		3
REEO PER	4	3	3	2 -		_	- (7 -		-	_	-	3	_		2		3	3	3	3		_		7
LARGE ANO ECGS ECGS	4	3	3	2 3		ı	5	7 -	2	2	4	2	2	2		2		-	2	-	-		4		_
S EGG	4	3	3	m r		2	5		1 ~	3	4	2	2	2		2		1	2	1	2		4		-
S LAYING	2	4	2	en er		1	- 4		2	3	3	3	3	2		3		4	3	1	2		,- -		-
SONIWORD &	4	4	3	ω 4		2	7	، ن	c 2	-	2	_	1	2		П		п	2	2	2		3		1
D AGE AT 8 50% PRO- 9 SUCTION	4	ĸ	3	2 4		2		٠ ر	1 -	-	Н	-	7	2		3		2	3	4	2		7	7	4
(Hew ponzed) O OUCTION EGG PRO-	4	4	3	т r		1	-	7 -		-	_	2	2	1		3		2	2	2	3		2		3
COST	4		- 3	- 5		- 1	1 (7 -			- 1	- 1	- 3	- 1				- 3		- 3	. 3		- 1		- 2
STRAIN OR TRADENAME	R.B.C.	1	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1			1	-300		3000-	- 1	-300	B-300	B-300	B-300		B-303		B-380	B-380	B-380	B-380				1
STRA	Kentville,	Anthony -		Anthony -		abcock	abcock	Babcock		abcock	Babcock	Babcock	Babcock	Babcock		Babcock		Babcock	Babcock	Babcock	Babcock		P. D. 58		Carey Nick 300
D NC	PS	SX	SX	x x		Z	Z i	Z 2	3 Z	Z	ZI	Z	Zī	Z		ZI		NBX	NBX	NBX	NBX		Syn.]		Z
BREEDING	WL	WL	WL	I M		WL	WL	M K		WL	WL	WL	WL	ML		WL		RIRXSYNBX	RIRXSYNBX	RIRXSYNBX	RIRXSYNBX		WL		WL
TEST	c, c.	MoC.	Mo. F.	Pa. Tenn		c.c.	Fla.	Minn.	Mo. F.	N.HC.	N.HF.	N.C.	Pa.	Tenn.		MoC.		MoF.	N. HC.	Z.C.	Pa.		C. C.		Minn.
ENTRY IDENTIFICATION	Animal Research Institute, Central Experimental Farm, Ottawa, Ontario, Canada.	Anthony, George M. & Sons, Strausstown, Pennsylvania 19559. Anthony, Pa		Anthony, Pa	Inc., P.O. Box 280, 350.	<pre>dt., Sask.)</pre>		1 1 1 1 1 1 1	N. Y. (Ballew, Mo.)		Babcock, N.Y.	Babcock, N.Y. (Beamsdale, N.C.)	Babcock, N.Y. (Babcock, Pa.)		Babcock Poultry Farm, Inc., P O. Box 280, Ithaca, New York 14850.	1 1 1 1 1 1 1	Babcock Poultry Farm, Inc., P.O. Box 280, Ithaca, New York 14850.	Babcock, N.Y	Babcock, N. Y.	Babcock, N.Y	Babcock, N.Y.	Canada Department of Agriculture, Poultry Division, Ottawa, Ontario, Canada.	Carev Farms, 3252 Mt. Olive. Agosta Road.		Carey, Ohio

TABLE 5. -- Range group ranking for stock entered in 1973-74 random sample egg production tests--Continued

ENTRY IOENTIFICATION	TEST	8.8	REEOING	STRAIN OR TRADENAME	OZT MO CHICK NEW LEED COWE	G PRO- ICTION (basuod na	% PRO - % DRO- 10T10N	VTIJATRO	YTIJATRO	THOIS	TRA LARGE 505 FEO PER	3G 2 00ND OE	YTIJAI	210
					10 S	H) 0	0S E	% GE			(3 E)	h (s)	10 (.)	ds (%)
Carey Farms, 3252 Mt. Olive, Agosta Road,														
1 1 1 1 1 1	MoF.	WL	Z	Nick	2	2	4	3	2	2	2	2	1	1
Carey, Ohio	Pa.	WL	Zī	Carey Nick 310	2	_	3	1		3	3	2	3	
Colonial Poultry Farm, Inc., Pleasant Hill,														
Missouri 64080.														
Colonial, Mo	Fla.	WL	Zī	True-Line 365 B-	ı	3	1	1		3	3	1	3	2
Colonial, Mo. (Ossenbrink, Mo.)	MoC.	WL	Zi	True-Line 365 B-	1	2	1	3	7	3	2	1	1	1
Colonial, Mo. (Research Farm, Mo.)	MoF	WL	Zi	True-Line 365 B-	2	4	4	4	3	4	3	1	1	1
Colonial, Mo	Pa,	WL	ZI	True-Line 365 B-	1	1	1	3	_	3	3	1	4	3
Colonial Poultry Farm, Inc., Pleasant Hill,														
Missouri 64080.														
Colonial, Mo	MoC.	WL	NI	True-Line 365 H-	1	3	2	3	2	3	3	1	1	1
Colonial Poultry Farm, Inc., Pleasant Hill,														
Missouri 64080.														
Colonial, Mo	MoC.	WL	Zi	True-Line 365 K-	1	3	1	2	2	3	3	1	1	1
Colonial, Mo. (Ossenbrink, Mo.)	MoF.	WL	Zi	True-Line 365 K-	2	2	2	4		3	3	1	1	1
Colonial Poultry Farm, Inc., Pleasant Hill,														
Missouri 64080.														
Colonial, Mo	C. C.	WL	ZI	True-Line 365 S.	2	3	2	1		3	3	Ī	3	2
Colonial, Mo	Minn.	WL	Z	True-Line 365 S-	2	2	1	1		3	3	2	3	3
1	MoF.	WL	ZI	True-Line 365 S-	1	2	2	3	_	3	3	1	I	1
Colonial, Mo	N. HC.	WL	Zī	True-Line 365 S-	3	3	7			4	4	2	3	3
Colonial, Mo	Tenn.	WL	ZI	True-Line 365 S-	2	3	3	4	2	3	3	1	3	_
Colonial Poultry Farm, Inc., Pleasant Hill,														
		1	ļ		,	,	(
', P.O. Box 27, Earl,	Mo. ¬F.	KIR	S.	Colonial RIR	4	4	~	~	4	m	m	4		
	0. F	RIRxBPR		70	4	4	£ (~		2		4	1 -	,
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	. HC.	RIRxBPR		το Ο	4	4	23	_		7	2	4	4	4
	H. F.	RIRxBF		avis	4	4	3	3	~	3	3	4	4	
Davis, N. C	Z.C.	RIR×BPR	R BX	Davis Combiner -	4	4	4	4		Ţ]	4	3	3

TABLE 5. -- Range group ranking for stock entered in 1973-74 random sample egg production tests--Continued

ENTRY IDENTIFICATION	TEST	88	BREEDING	STRAIN OR TRADENAME	CO21 ONEB CHICK ONEB LEED INCOME	S ONCLION ECC PRO-	B AGE AT SO OUCTION	YTIJATROM &	TIJATROM S	& WEIGHT	EGGS LARGE	(PS)	YTIJAUD S	STORS &
Euribrid B.V. Boxmeer, Holland Euribrid, Holland (Euribrid, Belgium)	MoC.	WL	SX			~4	6 0	2	4	2	2	2	1	
Euribrid, Holland	N. HC.	M K	XX S	White	7 -	7 -	ω -		، ٦	4 °	4 ~	7 -	m -	m r
Euribrid, Holland (Euribrid, Belgium) Fisher Poultry Farm, Ltd., Ayton, Ontario,	ر ر ر	ન ≱	Ś	nisex while	-	-	-	-	7	n	n	-	t	1
Canada.														
	C. C.	WL	SX		7	7	-	7	ش ا	7	7	ლ (4	7
Fisher, Ont	Minn.	WL	SX		ω.	m '	4 (ς (7 (ς,	7 (ς,	4	4
Fisher, Ont. (Bieman, Ont.)	MoC.	WL	XX S	Fisher 107	1 ~	4 ~	m r	2 "	7 ~	7 م	2 6	د د		1
Fisher, Ont.	MO. 15.] []	s k	107	. 4) 4	1 4	· ·) 4	1 m	1 ~	1 4	· (*)	_
Fisher, Ont.	Tenn.	W.E	SX	107	ر 2	7	, 2	2		2	- 2	2	4	2
Garber Poultry Breeding Farm, 4255 Hammett														
Koad, Modesto, California 79391.	<u>т</u>	WI	SX	Garber G 200		3	4	4	7	4	4	4	-	2
1	Minn.	WL	SX	Ü	3	4	3	2	3	4	4	4	1	-
Calif.	MoC.	WL	SX	Ü		3	3	3	7	3	3	3	,	,
Garber, Calif.	MoF.	WL	SX	Garber G 200	۴.	4	33	2	3	4	4	3	1	1
	Pa.	WL	SX	Garber G 200	. 5	3	2	4	4	4	4	7	2	_
Harco Farms, Bay Road, South Easton,														
Massachusetts 02375.												,		
Harco, Mass	Ħ.	-C.RIR×BPR		Sex	· ·	۳ ا	ر	ς,	٦.			4 (- .	- .
Harco, Mass	H.	-F. RIR×BPR	R BX	Harco Sex Link		7	m	_	ı	ı	-	2	_	_
Hardy, C. Nelson & Son, Essex,														
Massachusetts 01929.	1 1	T DID BDB	д Хд	Deluve Sev Link	~	~	4	_	2	2	-	~	3	3
Hubbard Farms, Inc., Walpole, New	:													
Hampshire 03608.	(;	·	~	-	<	_	0	~	4	4	~
Hubbard, N. H	N. H C. 2	SYNXN. H.	H. BX	i		n r	٦ ,	۰ -	7	1 -	, –	7 2	· ~	2
Hubbard, N.H		SYNXN. B.			1 m	1 W	ı ~		۱۳	- 2		4	~~	2
arms, P.O. Box 5														
Cameron, Texas 76520.											,	,		,
		SYN×WL				m i	7 ,	ლ (7 (٦,	7 .	7 °	ժ է	n -
	ا ب	SYN×WL			·	ν r	، ب	7 -	ν ~	n c	۰ ،	n -	, ,	۱.
Texas	MoF.	SYNXWL	X X	Ideal 236	7 -	7 0	v	۰ ر	o	7	٦ ر		. 4	٠ -
Ideal, lexas		SYN×W1	_		- -	ı —	, c	1 7	1 4	ı 2	2 2		4	. κ

TABLE 5. --Range group ranking for stock entered in 1973-74 random sample egg production tests--Continued

ENTRY IDENTIFICATION	TEST	BREEOING	٥ <mark>٧</mark>	STRAIN OR TRADENAME	COZT VAO CHICK OVER FEED INCOME	(Hew ponzed) S OUCTION EGG PRO-	DAGE AT SOUCTION	% GROWING % MORTALITY	VILLATROM %	& WEIGHT	LARGE ANO EGGS EGGS	C PEGGS (1PS)	YTIJAUQ 3	STORS &
Indiana Farm Bureau Coop., 2435 Kentucky Avenue, Indianapolis, Indiana 46204 Indiana Farm Bureau, Ind.	MoC.	M A	×	Duchess 60	'	4	2	4	41	~		23	1	
	MoF.	WL	SX	Duchess 60	2	2	2	3	2	4	. &	_	1	ı
1 1 1 1 1	Pa.	WL	SX	Duchess 60	1	_	2	-	3	4	4	2	1	_
1	Tenn.	WL	SX	Duchess 60	2	2	7	2	4	3	3	2	7	_
Kath Line of Canada, Box 415, Aldergrove, British Columbia, Canada														
	C. C.	WL	SX	Kath Line H 63	3	4	3	7	4	2	2	3	3	2
Lawton Farms, 70 North Street, Foxboro, Massachusetts 02035.														
	N.HF.RIR×WPR	IRxWPF	BX.	Buff Sex Link	4	4	4	4	_i	2		4	2	2
Missouri Valley Hatchery, Marshall, Missouri														
MolloV immo	Mo.	WI	×	Valley Oneen	3	۲.	4	2		~		2	1	
Lab.,		l :	;		b)		I))	i		
.7907.														
N.C. Reg. Plty., Ind N	Z.C.	WL	$_{\rm X}$	Kentville-Cornell	4	4	3	4	3	4	4	4	4	4
1 1 1 1 1 1 1 1	Tenn.	WL	$^{\rm SX}$	Kentville-Cornell	4	3	4	2	2	4	4	4	3	3
North Central Regional Poultry Breeding Lab.,														
N.C. Reg. Pltv Ind	Mo C.	WL	PS	Reg. Cornell Contr	1	4	4	3	3	4	4	4	1	1
1 1 1 1 1 1	Mo F.	WL	PS	Reg. Cornell Contr	. 3	3	4	3	2	4	4	3	\$	1
Parks Poultry Farm, Route 4, Box 118, Altoona, Pennsvlvania 16601.														
	Fla.	WL	SX	Keystone B-1	1	2	3	4	3		_	3	_	3
Parks, Pa N	Minn.	WL	SX	Keystone B-1	3	2	3	4	2	4	4	3	7	_
	N. HC.	WL	SX	Щ	3	3	3	1	3	4	4	3	<u>~</u>	
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Tenn.	WL	SX	Keystone B-1	3	3	IJ	4	2	4	4	3	2	3
Parks Poultry Farm, Route 4, Box 118,														
	MoF.RI	-F. RIR×WPR	BX	Sil-Go-Links	4	4	4	1	3	1	1	4	1	
Parks, Pa F	Pa. RI	RIR×WPR	BX	Sil-Go-Links	4	4	4		4	_	_	4	7	3

TABLE 5. -- Range group ranking for stock entered in 1973-74 random sample egg production tests--Continued

YTIJAUQ C. COOLITY	1 1 1 1 2 1 2 2 2 2 2 2 1 1 2 2 1 1 2 2 1 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 3 3 3 4 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5	5 - 1	11 - 2 - 4 - 2 - 4 - 4 - 4 - 4 - 4 - 4 - 4	4 -
LEECO PER CAND (%) ECCS (BS)	23 1 1 1 1 2 2 3 3 1 1 1 1 1 1 1 1 1 1 1	1 2 4		3 2 2
S EGG	2 3 2 2 2 5 1 1 2	L L 4		m m
SENDING (%) TILATROM (%) SUIYAL SUIYAL STILATROM (%)	3 3 1 1 3 3 1 1 1 1 1 1 1 1 1 1 1 1 1 1	3 2 2 1 2 3 4 4		1 2 2 1
D AGE AT	8 8 8 8 8 8 8	4.5 &	5 1 1 5 5 5 5 5 1 3	3 2
S ONCTION EGG PRO-	7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7	4.0 4	53535 11553	
CO21 PNO CHICK ONEE LEED INCOME		4.5 %		2 1
STRAIN OR TRADENAME	Starcross 288	Starcross 579 Starcross 579 Corvette A 1		Thornber 808 Thornber 808
BREEDING	$\overset{\circ}{\times}\overset{\circ}{\times}\overset{\circ}{\times}\overset{\circ}{\times}\overset{\circ}{\times}\overset{\circ}{\times}\overset{\circ}{\times}$	S X X	B B B B B B B B B B B B B B B B B B B	s× ××
в я	WL WL WL WL WL	RIR RIR WL	WL WL WL WL WL RIRXSYN RIRXSYN RIRXSYN RIRXSYN	WL
TEST	C.C. Fla. MoC. MoF. N.HC. N.C. Pa. Tenn.	C. C. MoF. C. C.	Fla Minn. MoF. Pa. Tenn. MoF. N.HC. Pa. Tenn.	C. C. MoC.
ENTRY IDENTIFICATION	Shaver Poultry Breeding Farm, Box 400, Galt-Cambridge, Ontario, Canada. Shaver, Ont.	ا أ ما أ	Jatum Farms, Koute 3, Dawsonville, Georgia 30534. Tatum, Ga	Thornber, England (Durance, Ont.) Thornber, England (Wilpstra, Ont.)

TABLE 5. -- Range group ranking for stock entered in 1973-74 random sample egg production tests--Continued

ENTRY IDENTIFICATION	TEST	B REE	BREEDING	STRAIN OR TRADENAME	CO21 VAD CHICK OAEK LEEO INCOME	Hem ponzeq) ECC PRO-	SO% PRO-	SEROWING TILLTY	SUIYAL STILITY	MEIGHT MEIGHT	ECCS	ECC2	YTIJAUD ;	STORS
Welp's Poultry Breeding Farm, Box 366,						1900	(ca)si	()	9	70	<u> </u>	(80)	4.0.7	Ŕ
Bancroft, Iowa 50517.														
	MoF.	RIR	SX	Welp Line 650 N-	3	2	_	1	3	2	~	3	,	,
1	N. HC.	RIR	SX	Welp Line 650 N-	3	2	2	-	П	2	2	4	3	7
Welp's Poultry Breeding Farm, Box 366,														
Bancroft, Iowa 50517.														
Welp, Iowa N	MoF.	WL	ZI	Welp Line 971	3	3	1	3	4	3	3	3	1	,
Welp, Iowa T	Tenn.	WL	ZI	Welp Line 971	3	3	2	3	3	3	3	23	3	2
Welp's Poultry Breeding Farm, Box 366,														
Bancroft, Iowa 50517.														
Welp, Iowa F	Fla.	WL	Z	Welp Line 973	1	4	3	4	4	3	3	4	4	4
Welp, Iowa N	MoF.	WL	ZI	Welp Line 973	3	4	2	4	4	3	3	3	ı	1
Welp, Iowa T	renn.	WL	ZI	Welp Line 973	4	4	1	2	4	3	3	4	4	_

RANDOM SAMPLE EGG PRODUCTION TEST ENTRIES AND CONDITIONS, 1973-74

TABLE 6. -- Stock entered in 1973-74 tests

× ×			X X X		× × × ×	×		oF. N. HC. N. HF. N. C. Pa. Tenn.	Tests entered
××	;			×					o × × ×
×	×				×	× ×			We
	×								X X X
			*	×	×	× ×	* *	× × ×	
	2 4	1 2 4	1 1 2 4	4 1 1 2 4	1 4 1 1 2 4	10 1 1 4 1 1 2 4 4 4 4 4 4 4 4 4 4 4 4 4 4	4 1 1 4 1 1 2 4	1 4 0 1 4 1 1 2 4	of entries 1 1 1 1 1 1 2 2 4 4 4 4 4 4 4 4 4 4 4 4
	365 B	10	B	B					1
Truelling	Carey Nick True-Line	Carey Nick 36 Carey Nick 31 True-Line 36	P. D. 58 Carey Nick 300 Carey Nick 310 True-Line 365	Babcock B-380 P. D. 58 Carey Nick 300 Carey Nick 310 True-Line 365	Babcock B-303 Babcock B-380 P. D. 58 Carey Nick 300 Carey Nick 310 True-Line 365	Babcock B-300 - Babcock B-380 - Babcock B-380 - P. D. 58 Carey Nick 300 - Carey Nick 310 - True-Line 365 E	Anthony Leghorn Babcock B-300 Babcock B-380 P. D. 58 Carey Nick 300 Carey Nick 310 True-Line 365 B		
our-en.I. 687								Kentville, R.B.C Anthony Leghorn- Babcock B-300 Babcock B-380 P.D. 58 Carey Nick 300	Kentville, R.B.C Anthony Leghorn- Babcock B-300 Babcock B-308 P.D. 58 Carey Nick 300

TABLE 6. -- Stock entered in 1973-74 tests--Continued

		Stock	Number					Test	Tests entered				
Breeder	Code	Strain or trade name	of entries	C. C.	Fla.	Minn.	MoC.	Mo F	N. H C.	N. HF.	Z, C	Pa.	Tenn.
Hubbard	378	Golden Comet	3						×		×	×	
Ideal	356	Ideal 236	Z.		×	×		×				×	×
Ind. Farm Bureau	234	Duchess 60	4				×	×				×	×
Kath	589	Kath-Line H-63	1	×									
Lawton	117	Buff Sex Link	1							×			
Missouri Valley	450	Valley Queen	1					×					
N. Cent. Reg. Lab	409	Kentville-Cornell	2								×		×
N. Cent. Reg. Lab	37	Reg. Cornell Control	2				×	×					
Parks	352	Parks Keystone B-1	4		×	×			×				×
Parks	382	Parks Sil-Go-Link	2					×				×	
Shaver	181	Shaver Starcross 288	∞	×	×		×	×	×		×	×	×
Shaver	451	Shaver Starcross 579	2	×				×					
St. Augustin	999	Corvette A-1	ı	×									
Tatum	401	Tatum T-100	Ŋ		×	×		×				×	×
Tatum	449	Tatum T-173	Ŋ					×	×	×		×	×
Thornber	407	Thornber 808	М	×			×	×					
Welp	440	Welp Line 650N	2					×	×				
Welp	430	Welp Line 971	2					×					×
Welp	448	Welp Line 973	3		×			×					×

TABLE 7. -- Management, rations, laying house environment, and vaccination provided by tests, 1973-74

			Length	Ent-	Replic	ations				Sq.
		Age at	of	ries		Birds	Hou	sing manager	nent	feet
Test	Hatched	housing (days)	test (days)	(num- ber)	Num- ber	per rep.	Brooding	Rearing	Laying 1/	per bird
Cent. Canada	3/27/73	147	497	12	4 4	65 65	Litter Litter	Litter Litter	Cage-2 Cage-2	0.45
Florida	6/ 3/73	150	486	12	4 8	24 75	Litter Litter	Litter Litter	Cage-2 Litter	.4 1.92
Minnesota	3/27/73	150	500	11 10	3 1	99 100	Litter Litter	Litter Litter	Cage-3 Litter	.33 1.5
Missouri Cage	9/ 9/72	151	500	14	2 4	40 40	Litter Litter	Litter Litter	Cage-2 Cage-8	.67 .58
Missouri Floor	3/ 3/73	151	500	27	4	60	Litter	Litter	Litter	1.6
New Hampshire Cage	5/ 1/73	150	502	17	8	24	Litter	Litter Cage	Cage-3	. 5
New Hampshire Floor	5/ 1/73	150	502	8	3	30	Litter	Litter	Litter	3.2
North Carolina	3/23/73	150	500	10	2 2	50 50	Litter Colony cage	Litter Colony cage	Litter-slat Colony cage-7	1.7 .5
					4	26	Colony cage	Colony	Cage-2	.6
Pennsylvania	4/23/73	150	500	24	2 2	48 50	Litter Litter	Litter Litter	Cage-3 Litter	.5 1.7
Tennessee	3/27/73	140	500	14	8	30	Litter	Litter	Cage-2	. 45

 $[\]underline{1}/$ The numerals after the word "cage" refer to the number of birds per cage.

TABLE 7. -- Management, rations, laying house environment, and vaccination provided by tests, 1973-74

Continued

Test	Entries brooded inter-	Min. oz./doz. for large		Protein (percent)			energy ies/pound		MC/Ci	· Prot.	3/
	mingled	eggs	Start	Grow	Lay	Start	Grow	Lay	Start	Grow	Lay
Cent. Canada - Test Control Ration		24	14.7 20.3	10.5 16.2	16.9 16.9	1270 1270	1290 1290	1300 1300	80.9 58.0	124.0 79.1	
Florida	Yes	23	22.0	9.1 15.3	16.5	1340	1480 1371	1313	60.9	162.6 78.8	77.7
Minnesota	Yes	23	20.2	15.2	17.5	1268	1215	1310	63.0	80.0	74.9
Missouri Cage	No	23	20.7	16.2	18.2 15.1	1318	1266	1250 1224	63.7	78.1	68.7 81.1
Missouri Floor	No	23	20.7	16.2	17.0 15.1	1318	1266	1281 1305	63.7	78.1	75.3 86.4
New Hampshire	Yes	23.5	20.9	16.0	18.5 to 15.5	1340	1319	1255 to 1337	64.0	82.0	72.0 to 81.0
North Carolina	No	23	20.0	16.0	18.0 to 16.0	1249	1238	1303 to 1335	62.4	77.4	71.2 to 80.9
Pennsylvania	Yes	24	21.0	17.0	18.0	13004/	13574/	1354 <u>4</u>	61.9	79.8	75.2
Tennessee	- No	23	20.8	16.5 9.0	$16.9\frac{5}{5}$ 16.9	/ / 1365 / 1365	1382 1443	1305 1305	65.6 65.6	84.0 1 5 9.0	

^{2/} Metabolizable energy is the maximum quantity of feed energy that possibly may be used by the chicken.

^{3/} Metabolizable calories divided by percent crude protein.

^{4/} Approximate metabolizable energy computed from productive energy, using 70 percent as the conversion factor.

^{5/} See Tennessee Test Report for complete ration combinations.

TABLE 7. -- Management, rations, laying house environment, and vaccination provided by tests, 1973-74

Continued

	Ligl	nting	Artificial			
Test	Rearing (hours)	Laying (hours)	heat used	R Value of insu material <u>6</u> /		Ventilation
Cent. Canada	(<u>7</u> /)	(<u>8</u> /)	Yes	Ceiling Walls	27.9 15.1	Exhaust fans in roof and in east wall.
Florida	Natural	15	No	Cage Summer House Winter	13.0 8.0	Natural ridge vents
Minnesota Cage	12	12 to 16	No	Ceiling Walls	15.8 12.1	Positive pressure
Minnesota Floor	Natural	12 to 16	No	Ceiling Walls	15.0 13.0	Exhaust fans
Missouri Cage	10	16	No	Ceiling Walls	5.8 None	Ridge vents
Missouri Floor	Natural	14	No	Ceiling Walls	15.0 15.0	Exhaust fans in ceiling
New Hampshire	14	14	No	Ceiling Walls	15.0 15.0	Exhaust fans
North Carolina	Step down	Step up to 17	No	Ceiling Walls	7.3 1.5	Natural via windows
Pennsylvania	8	12 to 17	Yes	Ceiling Walls	15.5 15.5	Exhaust
Tennessee	Natural	14	No	Ceiling Walls	13.0 None	Winter, Positive pressure; Summer, Exhaust fans

^{6/} Due to variations in type of construction, R Values will be approximate for some tests.

^{7/} At day old--18-1/2 hr.; light decreased 15 minutes per wk. to meet at 15-1/2 hr. at longest day, then natural decrease until 13-1/2 hr.

^{8/ 13-1/2} hr. until natural increase takes light hours to 15-1/2 hr. in mid-June, then light held at 15-1/2 hr. until end of test.

TABLE 7.-- Management, rations, laying house environment, and vaccination provided by tests, 1973-74

Continued

	Newca	stle	Infect broncl		Fowl	Pox		phalo- elitis	Coccidiosi control		Marek's Disease
Test	Туре	Age (wk.)	Туре	Age (wk.)	Туре	Age (wk.)	Туре	Age (wk.)	Туре	Age (wk.)	Age
Central Canada	Spray Spray	1.5 19	Spray Spray	1.5 12	Wing web.	8	Water Water	8 15			l day
Florida	Water Water Water	1,3,10 16,32 48,64	Water Water	1,3 10,16	Wing web.	8	None		Poly-stat	0-8	l day
Minnesota	Water Water	5 14	Water Water	5 14	Wing web.	9	None		Amprol	0-20	l day
Missouri Cage	Water Water Water	1 7 16	Water Water Water	1 7 16	None		None		Poly-stat	0-11	l day
Missouri Floor	Water Water Water	1 4 10	Water Water Water	1 4 10	None		None		Poly-stat	0-8	1 day
New Hamp- shire	- Dust Dust	2 20	Dust Dust	2 20	None		None		Cocci-Vac	1	l day
North Carolina	Occular Water Water +Every 90	1 5 16) days	Occular Water	1 5	Wing web.	12	Water	14	None (cages) 6 Spcs. Cocc		l day
Pennsyl- vania	Water Water Water	4 8 16	Water Water Water	4 8 16	Wing web.	8	None		Amprol	0-20	l day
Tennessee	Occular Occular Occular	1 day 10 20	Occular Occular Occular	1 day 10 20	Wing web.	10	None		Amprol	0 -20	l day

GPO 882-991

U. S. DEPARTMENT OF AGRICULTURE AGRICULTURAL RESEARCH SERVICE NORTHEASTERN REGION AGRICULTURAL RESEARCH CENTER WEST BELTSVILLE, MARYLAND 20705

OFFICIAL BUSINESS
PENALTY FOR PRIVATE USE, \$300

POSTAGE AND FEES PAID
U. S. DEPARTMENT OF
AGRICULTURE
AGR 101

